Edward Grzegorek A remark on a paper by Karel Prikry

In: Zdeněk Frolík (ed.): Abstracta. 5th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1977. pp. 23–25.

Persistent URL: http://dml.cz/dmlcz/701081

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

FIFTH WINTER SCHOOL (1977)

A REMARK ON A PAPER BY KAREL PRIKRY

ру

E. GRZEGOREK

Our note is a remark on a paper [2] by Prikry. In the present note, an ordinal is considered to be the set of the smaller ordinals. Cardinals are the initial ordinals. Moreover, we shall adopt the convention that ξ and φ denote ordinals, and that \varkappa and λ denote infinite cardinals. If S is a set then $\mathfrak{P}(S)$ denotes the family of all subsets of S. If $\mathcal A$ is a field of subsets of S, then by $\mathcal J(\mathcal A)$ we denote the family of all A&A such that for every XCA we have X $\in \mathcal{A}$. A family of sets \mathfrak{R} is said to be \varkappa - complete if for each family $\mathfrak{F} = \{A_{\xi}: \xi < \lambda\} \subset \mathfrak{R}$, where $\lambda < \mathcal{H}$, $U\mathcal{F} \in \mathcal{R}$ as well. Let \mathcal{J} be an ideal in \mathcal{P} (S). A family \mathcal{R} C \mathcal{P} (S) satisfies (x.C.C.)(\mathcal{J}) iff each family \mathcal{F} C \mathcal{R} such that $F \cap G \in \mathcal{J}$ for all $F,G \in \mathcal{F}$, $F \neq G$ has power $< \mathcal{H}$. We always assume $\emptyset \in \mathcal{J}$. If $\mathcal{J} = \{\emptyset\}$, then instead of (X.C.C.)(J) we simply write X.C.C. The almost disjoint transversals hypothesis for ω_1 is denoted by TH(ω_1). Let us recall that TH(ω_1) follows from Gödels axiom of constructability (for more information see [2]).

By modifications of Prikry's proofs and definitions in [2] we generalize theorems of [2] to the following form.

Theorem 1. Assume TH(ω_1). Let \mathcal{M} be a family of ω_1 -complete fields of subsets of ω_1 such that $|\mathcal{M}| \leq \omega_1$ and for every $\mathcal{A} \in \mathcal{M}$ the family $\mathcal{A} - J(\mathcal{A})$ satisfies ω_1 .c.c. and $UJ(\mathcal{A}) = \omega_1$.

Then for every $X \notin \bigcup \{J(\mathcal{X}) : \mathcal{X} \in \mathcal{M}\}$ there exists a disjoint family $\{X_{\xi} : \xi \in \omega_1\} \subset \mathcal{P}(X)$ such that $X_{\xi} \notin \mathcal{H}$ for every $\mathcal{A} \in \mathcal{M}$ and every $\xi \in \omega_1$.

Theorem 2. Let \mathcal{H} and λ be regular cardinals such that $\lambda \geqslant \mathcal{H} \geqslant \omega_1$ and let \mathcal{M} be a countable family of \mathcal{H} -complete fields of subsets of S such that for every $\mathcal{H} \in \mathcal{M}$ the family $\mathcal{H} - J(\mathcal{H})$ satisfies $\mathcal{H} \cdot C.C.$. Suppose $J \subset \bigcap \{J(\mathcal{H}): \mathcal{H} \in \mathcal{M}\}$ is a \mathcal{H} -complete ideal in $\mathcal{H} \cap \mathcal{H}$ (S) such that for every $\mathcal{H} \in \mathcal{M}$ and every $\mathcal{H} \in \mathcal{H}$ the family $\mathcal{H} \cap \mathcal{H}$ (S) and every $\mathcal{H} \in \mathcal{H}$ (S) $\mathcal{H} \cap \mathcal{H}$ the family $\mathcal{H} \cap \mathcal{H}$ (S) there exists a family $\mathcal{H} \cap \mathcal{H}$ (S) such that

exists a family $\{X_{\xi}: \xi \in \lambda\} \subset \mathcal{P}(X)$ such that

(1) $X_{\xi} \cap X_{\varphi} \in \mathcal{J}$ for every $\xi \neq \varphi$, $\xi, \varphi \in \lambda$, and

(11) for every $\xi \in \lambda$, every $\mathcal{H} \in \mathcal{M}$ and every $A \in \mathcal{H}$ we have, if $X_{\xi} - A \in \mathcal{J}(\mathcal{H})$, then $X - A \in \mathcal{J}(\mathcal{H})$.

Theorem 3. Let λ be a regular cardinal and let $\mathcal M$ be a finite family of fields of subsets of S such that for every $\mathcal K \in \mathcal M$ the family $\mathcal K = \mathcal J(\mathcal K)$ satisfies $\omega_0.\text{C.C.}$. Suppose $\mathcal J \in \mathcal J(\mathcal K): \mathcal K \in \mathcal M$ is an ideal in $\mathcal J \in \mathcal J(\mathcal K)$ such that for every $\mathcal K \in \mathcal J$ and every $\mathcal K \in \mathcal J(\mathcal K) = \mathcal J(\mathcal K)$ the family $\mathcal J \in \mathcal J(\mathcal K) = \mathcal J(\mathcal K)$ does not satisfy $\{\mathcal K \in \mathcal J \cap \mathcal J \in \mathcal J \cap \mathcal$

In [2] Prikry has proved theorems 1 and 2 in the case if \mathcal{H} consists only ω_1 - complete fields on which it is possible to define complete probabilities which vanishe on all finite sets. Theorem 1 of this note generalizes Theorem 1 of [2], and Theorem 2 of this note generalizes Theorem 2 of [2]. Our generalization of Theorem 2 of [2] may be motivated by the fact that it gives a common generalization and

strenghtening of known theorems for measures (see Sierpiński [4] and Prikry [2]), for outer measures (see Poprużenko [1]), and for the category of Baire (see Sierpiński [3]).

An extended form of this note will submitted for publication elsewhere.

References

- [1] J. Popruženko, Sur une propriété d'une classe de measures abstraites, Colloquium Mathematicum 4(1957), p. 189-194.
- [2] Karel Prikry, Kurepa's hypothesis and a problem of Ulam on families of measures, Monatshefte für Mathematik 81, (1976), p. 41-57.
- [3] W. Sierpiński, Sur les ensembles partout de deuxième catégorie, Fundamenta Mathematicae 22(1934), p. 1-3.
- [4] -, Sur une propriété des ensembles linéaires quelconques, Fundamenta Mathematicae 23(1934), p. 125-134.