

Zbigniew Lipecki

Extension of functions on Boolean algebras

In: Zdeněk Frolík (ed.): Abstracta. 5th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1977. pp. 39–42.

Persistent URL: <http://dml.cz/dmlcz/701087>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

EXTENSION OF FUNCTIONS ON BOOLEAN ALGEBRAS

BY

Z. LIPECKI

Let A be a Boolean algebra with the operations of join, meet and difference denoted by $\vee$, $\wedge$ and $\setminus$, respectively. The natural ordering of A is denoted by $\leq$ and its minimal element by 0 .

Given a subalgebra B of A and a function μ on B with values in $[0, \infty)$ or in an Abelian (topological) group G which has some properties related to the structures of both the domain and range spaces, we consider the problem of extending μ to A all the properties preserved.

I. Suppose $\mu: B \rightarrow [0, \infty)$. We are concerned with four sets of conditions imposed on μ .

(i) μ is monotone (i.e. $\mu(b) \leq \mu(c)$ whenever $b, c \in B$ and $b \leq c$), subadditive (i.e. $\mu(b_1 \vee b_2) \leq \mu(b_1) + \mu(b_2)$ whenever $b_1, b_2 \in B$) and $\mu(0) = 0$ (').

(ii) μ is additive (i.e. $\mu(b_1 \vee b_2) = \mu(b_1) + \mu(b_2)$ whenever $b_1, b_2 \in B$ and $b_1 \wedge b_2 = 0$).

(iii) μ is monotone, subadditive and exhaustive (i.e. $\mu(b_i) \rightarrow 0$ whenever $\{b_i\} \subset B$ and $b_i \wedge b_{i'} = 0$ for all $i \neq i'$).

(iv) μ is monotone, exhaustive and $\mu(b \vee c) = \mu(b \setminus c)$ ($= \mu(b)$) whenever $b, c \in B$ and $\mu(c) = 0$.

(') Such a function is usually called a submeasure.

As easily seen, $(ii) \Rightarrow (i)$ and $(ii) \Rightarrow (iii) \Rightarrow (iv)$ and none of these implications can be reversed.

The case where μ satisfies (i) is trivial. Putting $\nu(a) = \inf \{\mu(b) : a \leq b \in B\}$ for $a \in A$, we get an extension satisfying (i).

The answer is also positive in case μ fulfils (ii) or (iii). As for (ii), this is a classical result of Hahn-Banach type (see, e.g., [4], p. 270). When dealing with (iii), the transfinite argument must be essentially improved since the exhaustivity condition involves countably many elements of A . The improvement is suggested by the following observations:

1. For any μ satisfying (i) the formula $d_\mu(b_1, b_2) = \mu((b_1 \vee b_2) \vee (b_1 \wedge b_2))$ for $b_1, b_2 \in B$ defines a pseudometric on B .
2. If $C \subset B$ is a d_μ -dense subalgebra of B and $\mu|_C$ is exhaustive, then so is μ .

(The details can be reconstructed from [3]. For a different proof see [2].)

As for (iv), the answer is positive under the additional assumption that B has the following compactness type property investigated by Seever [5]:

- (I) Suppose $b_k, c_k \in B$ and $\bigvee_{i=1}^n b_i \leq \bigvee_{i=1}^n c_i$ for $k=1, 2, \dots$. Then there exists $x \in B$ with $\bigvee_{i=1}^n b_i \leq x \leq \bigvee_{i=1}^n c_i$ for $k=1, 2, \dots$.

This property, which is somewhat weaker than σ -completeness, corresponds to the fact that the representation space of B is an F -space ([5], Theorem A). The author does not know whether the result fails for B without (I) ⁽²⁾. Before sketching

⁽²⁾ For an arbitrary B it can be shown that if μ fulfils (iv) with "exhaustive" weakened to " $\mu(0) = 0$ ", then it extends to A all its properties preserved.

the proof we give a lemma which is a slight generalization of a result due to Seever ([5], Lemma 3.3).

Lemma. If B has the property (I), N is an ideal in B and B/N satisfies the countable chain condition (CCC), then B/N is complete.

Suppose μ satisfies (iv) and B has the property (I). Put $N = \{b \in B : \mu(b) = 0\}$. Then B/N satisfies CCC, so that, by the Lemma and Sikorski's theorem ([6], 33.1), the quotient homomorphism $B \rightarrow B/N$ extends to a homomorphism $h: A \rightarrow B/N$. Putting $\nu(a) = \bar{\mu}(h(a))$, where $\bar{\mu}: B/N \rightarrow [0, \infty)$ is the quotient of μ , we get the desired extension (cf. also [2], Theorem 1).

II. Suppose G is an Abelian complete Hausdorff topological group and $\mu: B \rightarrow G$ is additive and exhaustive. (These properties are defined in just the same way as in I.) Then μ extends to A both properties preserved ([3], Theorem 3, and [2], Corollary 3). This theorem can be proved using a similar idea to that described when dealing with μ satisfying (iii). In fact, it is not hard to see that the group-valued case is more general than (iii). Let us also note that the assumption that G is complete cannot be dropped ([3], Example 4).

III. Suppose G is an Abelian group and $\mu: B \rightarrow G$ is additive. In this case the extension problem is open in general (e.g. for G being the additive group of integers). The answer is positive under each of the following additional assumptions:

(a) (W. Herer) G is injective (see [11], §21, for definition).

(b) (C. Ryll-Nardzewski) $\text{card } A \leq \aleph_1$.

(c) B/N is complete, where $N = \{b \in B : \mu(c) = 0 \text{ for all } b \geq c \in B\}$.

As for (c), the assertion follows by an application of Sikorski's theorem analogously to the case where μ satisfies (iv).

REFERENCES

- [1] L. Fuchs, Infinite Abelian Groups, Vol. I, Academic Press, 1970.
- [2] S. S. Khurana, Extensions of exhaustive submeasures, Bull. Acad. Polon. Sci., Série Sci. Math. Astr. Phys. 24 (1976), p. 213-216.
- [3] Z. Lipecki, Extensions of additive set functions with values in a topological group, ibidem 22 (1974), p. 19-27.
- [4] J. Łoś and E. Marczewski, Extensions of measure, Fund. Math. 36 (1949), p. 267-276.
- [5] G. L. Seever, Measures on F-spaces, Trans. Amer. Math. Soc. 133 (1968), p. 267-280.
- [6] R. Sikorski, Boolean Algebras, Springer-Verlag, 1964.