

Erik G. F. Thomas

A Krein-Milman set without the integral representation property

In: Zdeněk Frolík (ed.): Abstracta. 8th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1980. pp. 39–41.

Persistent URL: <http://dml.cz/dmlcz/701173>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1980

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

A KREIN - MILMAN SET WITHOUT THE INTEGRAL REPRESENTATIONPROPERTY

by

Křesomysl Blizzard

We construct a separable Banach space E and a bounded, closed, absolutely convex subset B such that B is the closed convex hull of its extreme points but such that not every point in B is representable as the barycenter of a probability measure on the extreme points of B .

Let X be a separable Banach space not having the Radon-Nikodym property and such that its unit ball U is the closed convex hull of its extreme points $E(U)$. The space of converging sequences c for example is such a space. (Note in passing that the unit ball of c has countably many extreme points and that every point in the unit-ball of x is the barycenter of probability measure on the extreme points).

Let A be the Cantor set and let $E = I(C(A), X)$ be the space of integral operators from $C(A)$ to X , i.e. the linear operators $T : C(A) \rightarrow X$ such that

$\|T\|_I = \sup \{ \sum_{i=1}^n \|T \chi_{A_i}\| : A_i \text{ disjoint clopen sets in } C \} < \infty$.
Let $B = \{T : \|T\|_I \leq 1\}$ and equip E with the topology τ of pointwise convergence on $C(A)$, i.e. $T_\alpha \rightarrow T$ if for each $f \in C(A)$, $\|T_\alpha(f) - T(f)\| \rightarrow 0$.

There are obvious extreme points in B , namely the $\delta_t \otimes x_f$ $t \in A$, $x \in E(U)$. It is also obvious that these are the only extreme points of B , hence we write

$E(B) = \{\delta_t \otimes x, t \in \Delta, x \in E(U)\}$. We shall show that B is the closed convex hull of $E(B)$.

By the Hahn-Banach theorem this is equivalent to say that the polars of $E(B)$ and B coincide. Let

$\sum_{i=1}^n f_i \otimes x_i^*$ be an element of E' , that belongs to the polar of $E(B)$. Evidently this means just that for $t \in \Delta$,

$\|\sum_{i=1}^n f_i(t) \cdot x_i^*\|_{X^*} \leq 1$ and this latter condition implies that

$\sum_{i=1}^n f_i \otimes x_i^*$ belongs to the polar of B , as is readily seen from the definition of B . Hence $\bar{E}(B) = B$.

We shall now show that there are points in B not representable as barycenter of probability measures on the extremals. Let T_0 be an integral operator in B that is not nuclear and suppose there is a probability μ on $E(B)$ such that for each $f \in C(\Delta)$ and $x^* \in X^*$

$$\langle x^*, T_0(f) \rangle = \int_{E(B)} \langle x^*, \delta_t \otimes x \rangle d\mu(\delta_t \otimes x).$$

Note that $E(B)$ is homeomorphic to $E(U) \times \Delta$. As $E(U)$ is always a coanalytic set (if X is c , $E(U)$ is even a countable discrete set), there exists a desintegration of μ , i.e. there are probability measures μ_t on E_U and a probability measure ν on Δ such that $\mu = \int_{\Delta} \mu_t d\nu(t)$, i.e. we get for $f \in C(\Delta)$ and $x^* \in X^*$

$$\begin{aligned} \langle x^*, T_0(f) \rangle &= \int_{\Delta} \left[\int_{E_U} \langle x^*, (\delta_t \otimes x)(f) \rangle d\mu_t(x) \right] d\nu(t), \\ &= \int_{\Delta} f(t) \cdot \left[\int_{E_U} \langle x^*, x \rangle d\mu_t(x) \right] d\nu(t). \end{aligned}$$

For $t \in \Delta$ write $F(t) = \int_{E_U} x d\mu_t(x) \in U$ (the integral taken in the weak sense) to get a Radon-Nikodym derivative of T_0 , i.e. for $f \in C(\Delta)$

$$T_0(f) = \int_{\Delta} f(t) \cdot F(t) d\nu(t).$$

This means just that T_0 is nuclear, which is a contradiction.

Remark: We have constructed our example in a locally convex space E which is not even a Fréchet space, but it is not difficult to make the example live in a Banach space. Let $\{f_n\}_{n=1}^\infty$ be any total sequence in $C(\Delta)$, tending to zero in norm, and define the norm $\|\cdot\|_E$ in E to be

$$\|T\|_E = \sup \{ \|T f_n\| : n \in \mathbb{N} \}.$$

It is easily verified that $\|\cdot\|_E$ is indeed a norm and defines on B the topology τ . Letting $\tilde{E}$ be the completion of $(E, \|\cdot\|_E)$ we have imbedded our example into a separable Banach space.

An inspection of the above argument shows, that we may imbed our example into the space $c_0(X)$ or even $l^2(X)$.