

Martin Gavalec

Products of ideals of Borel sets

In: Zdeněk Frolík (ed.): Abstracta. 9th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1981. pp. 32–33.

Persistent URL: <http://dml.cz/dmlcz/701222>

## Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1981

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

## PRODUCTS OF IDEALS OF BOREL SETS

Martin Gavalet

The natural definition of the product  $\mathcal{I} \times \mathcal{J}$  of ideals in the fields  $\mathcal{B}(X)$ ,  $\mathcal{B}(Y)$  of all Borel sets in topological spaces  $X, Y$  sounds as follows:

for any  $A \in \mathcal{B}(X \times Y)$  we set

$$A \notin \mathcal{I} \times \mathcal{J} \equiv \{x \in X; \{y \in Y; (x, y) \in A\} \notin \mathcal{J}\} \notin \mathcal{I}.$$

This definition is meaningful if

(\*) the sets  $\{y \in Y; (x, y) \in A\}$  for  $x \in X$  are Borel in  $Y$ , and if

(\*\*) the set  $\{x \in X; \{y \in Y; (x, y) \in A\} \notin \mathcal{J}\}$  is Borel in  $X$ .

The first condition is always satisfied, the second one depends on the ideal  $\mathcal{J}$ .

Let us denote by  $\mathcal{L}, \mathcal{K}$  the ideals of all Borel sets in the real unit interval  $I$ , of the Lebesgue measure zero, or of the first Baire category, respectively.

Theorem 1. If the ideal  $\mathcal{J}$  is a product of finitely many ideals, each equal to  $\mathcal{L}$ , or to  $\mathcal{K}$ , then the condition (\*\*) is satisfied.

Theorem 1 enables us to form products of ideals  $\mathcal{L}, \mathcal{K}$ , in arbitrary order. The following theorem describes an important property of such products.

Theorem 2. If the ideal  $\mathcal{J}$  is the product of  $m$  ideals, each equal to  $\mathcal{L}$  or to  $\mathcal{K}$ , then  $\mathcal{J}$  is countably complete and the boolean algebra  $\mathcal{B}(I^m)/\mathcal{J}$  fulfills the countable chain condition. The algebra  $\mathcal{B}(I^m)/\mathcal{J}$  is, therefore, complete.

Complete boolean algebras and their complete boolean products are closely connected with boolean-valued models of the axiomatic set theory. In [1], [2] the property of local disjointness is described, which is fulfilled in a complete boolean product if and only if the corresponding model classes are disjoint over the basic model.

Theorem 3. If the ideal  $\gamma$  is the product of  $m$  ideals,  $k$  of which (not necessarily the first ones) are equal to  $\mathbb{L}$  and  $m-k$  are equal to  $\mathbb{K}$ ,  $0 < k < m$ , then the complete product  $\mathcal{B}(\mathbb{I}^m)/\gamma$  of algebras  $\mathcal{B}(\mathbb{I}^k)/\mathbb{L}^k$ ,  $\mathcal{B}(\mathbb{I}^{m-k})/\mathbb{K}^{m-k}$  induced by the natural embeddings, is locally disjoint.

Remarks. 1. By the well-known Fubini's theorem, the algebra  $\mathcal{B}(\mathbb{I}^k)/\mathbb{L}^k$  is isomorphic to the so-called random algebra  $\mathcal{R} = \mathcal{B}(\mathbb{I})/\mathbb{L}$ . Analogously,  $\mathcal{B}(\mathbb{I}^{m-k})/\mathbb{K}^{m-k}$  is isomorphic to the Cantor algebra  $\mathcal{C} = \mathcal{B}(\mathbb{I})/\mathbb{K}$ . Thus, Theorem 3 is a tool for constructing infinitely many non-isomorphic locally disjoint products of algebras  $\mathcal{R}, \mathcal{C}$ .

2. The product of algebras  $\mathcal{R}, \mathcal{C}$ , described above are non-isomorphic when considered as products. It is a problem, if they are isomorphic as boolean algebras. E.g. are the boolean algebras  $\mathcal{B}(\mathbb{I}^2)/\mathbb{L} \times \mathbb{K}$ ,  $\mathcal{B}(\mathbb{I}^2)/\mathbb{K} \times \mathbb{L}$  isomorphic?

#### References

- [1] L. Bukovský, Cogeneric extensions, Proc. Wrocław Logic Coll. 1977, North Holland 1978, 91-98.
- [2] M. Gavalec, Local properties of complete boolean products, to appear in Coll. Math.