

Zhen Yuan Xu

On boundary value problem of Neumann type for hypercomplex function with values in a Clifford algebra

In: Jarolím Bureš and Vladimír Souček (eds.): Proceedings of the Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1990. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 22. pp. [213]–226.

Persistent URL: <http://dml.cz/dmlcz/701474>

### Terms of use:

© Circolo Matematico di Palermo, 1990

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

# ON BOUNDARY VALUE PROBLEM OF NEUMANN TYPE FOR HYPERCOMPLEX FUNCTION WITH VALUES IN A CLIFFORD ALGEBRA

Xu Zhenyuan

## 1. Introduction

Let  $R_{0,n-1}$  be the universal algebra constructed by means of an orthonormal basis  $\{e_2, e_3, \dots, e_n\}$  of  $R^{0,n-1}$  and  $e_\phi = e_1$  be the identity of  $R_{0,n-1}$ . In the universal Clifford algebra  $R_{0,n-1}$  the multiplicative identity

$$e_i e_j + e_j e_i = -2\delta_{ij} \quad i, j = 2, 3, \dots, n,$$

holds. Let  $\bar{\partial} = e_1 \frac{\partial}{\partial x_1} + \sum_{j=2}^n e_j \frac{\partial}{\partial x_j}$  be a generalized Cauchy-Riemann operator

and  $\Omega$  be an open set of  $R^n$ . Then an  $R_{0,n-1}$ -valued  $C^1(\Omega)$ -function  $f(x) =$

$\sum_A f_A(x) e_A$  is called left monogenic function if  $\bar{\partial} f = 0$  in  $\Omega$ , here  $e_A =$

$e_{\alpha_2 \alpha_3 \dots \alpha_h} = e_{\alpha_2} e_{\alpha_3} \dots e_{\alpha_h}$  are bases of  $R_{0,n-1}$  and  $A = \{\alpha_2, \alpha_3, \dots, \alpha_h\} \subseteq \{2, 3, \dots, n\}$  with  $2 \leq \alpha_2 < \dots < \alpha_h \leq n$ .

Up to now a great number of the function theory for left monogenic function has been investigated [1][2][6]. For the use of Clifford algebra in several branches of Mathematics and Physics we refer to [2]. Recently several boundary value problem for monogenic functions have been studied (see [8][9][10][11]). In this paper we study a boundary value problem of Neumann type for hypercomplex function with values in a Clifford algebra which is a solution of a nonhomogeneous equation  $\bar{\partial} w = f$  in the

---

This paper is in final form and no version of it will be submitted for publication elsewhere.

unit ball of  $\mathbb{R}^n$ . To this end we firstly construct a Neumann's function for Laplacian over the unit ball in  $\mathbb{R}^n (n \geq 3)$ . Then, making use of this Neumann's function, we solve the above boundary value problem.

Acknowledgment This paper was written during the author's stay at Gent, which was supported by State University of Gent scholarship. She wishes to express her thanks to Prof. Dr. Delanghe for his support.

## 2. Neumann's function

In this section we construct an explicit representation formula for Neumann's function for Laplacian over the unit ball in  $\mathbb{R}^n (n \geq 3)$  by means of the fundamental solution for Laplacian and Gegenbauer polynomials. Then, making use of this Neumann's function, the solution of the Neumann problem for the Poisson equation is represented.

In the sequel,  $K_n$  denotes a unit ball in  $\mathbb{R}^n$ ,  $K_n = \{x \mid x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n, |x| < 1\}$ . We seek for a real function  $N(x, y)$  which satisfies

$$\Delta_x N(x, y) = \delta(x - y) + c, \quad x \in K_n \quad (2.1)$$

$$\frac{\partial}{\partial n_x} N(x, y) = 0, \quad x \in \partial K_n \quad (2.2)$$

hereby  $\Delta_x$  denotes the Laplacian  $\Delta_x = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$ ,  $\frac{\partial}{\partial n_x}$  denotes the outer normal derivative to the boundary  $\partial K_n$  of  $K_n$ ,  $c$  is a suitable constant to be determined, while  $y$  is fixed in the domain  $K_n$ . In fact this problem is not solvable for any real constant  $c$ .

Now we try to look for a solution of problem (2.1)(2.2) in the following form

$$N(x, y) = \frac{1}{\omega_n(2-n)|x-y|^{n-2}} + \frac{1}{\omega_n(2-n)|y|^{n-2}|x-y^*|^{n-2}} + g(x, y), \quad (2.3)$$

whereby  $\omega_n$  stands for the area of the unit ball in  $\mathbb{R}^n$ ,  $\omega_n = 2\pi^{n/2}/\Gamma(n/2)$ ,  $\frac{1}{\omega_n(2-n)|x-y|^{n-2}}$  is the fundamental solution for Laplacian, namely

$\Delta_x \left( \frac{1}{\omega_n(2-n)|x-y|^{n-2}} \right) = \delta(x-y)$ ;  $y^*$  is the symmetric point of  $y$  to the unit

sphere  $\partial K_n$ ,  $y^* = \frac{y}{|y|^2}$ . It is clear that, for  $y \in K_n$ , the function  $\frac{1}{\omega_n(2-n)|y|^{n-2}|x-y^*|^{n-2}}$  is harmonic in  $K_n$  with respect to  $x$ .

By (2.1)(2.2), for  $y \in K_n$ , the function  $g(x, y)$  should satisfies

$$\Delta_x g(x, y) = c, \quad x \in K_n \quad (2.4)$$

$$\frac{\partial}{\partial n_x} g(x, y) = - \frac{\partial}{\partial n_x} \left( \frac{1}{\omega_n(2-n)|x-y|^{n-2}} + \frac{1}{\omega_n(2-n)|y|^{n-2}|x-y^*|^{n-2}} \right), \quad x \in \partial K_n \quad (2.5)$$

Now we suppose that the function  $g(x, y)$  has the following form

$$g(x, y) = h(x, y) + \alpha |x|^2, \quad (2.6)$$

so that the function  $h(x, y)$  satisfies, for  $y \in K_n$ ,

$$\Delta_x h(x, y) = 0, \quad x \in K_n \quad (2.7)$$

$$\frac{\partial}{\partial n_x} h(x, y) = - \frac{\partial}{\partial n_x} \left( \frac{1}{\omega_n(2-n)|x-y|^{n-2}} + \frac{1}{\omega_n(2-n)|y|^{n-2}|x-y^*|^{n-2}} + \alpha |x|^2 \right), \quad x \in \partial K_n \quad (2.8)$$

Thus the constants  $c$  and  $\alpha$  are related with

$$c = 2n\alpha. \quad (2.9)$$

It is well known that the Neumann problem (2.7)(2.8) is solvable if and only if the constant  $\alpha$  satisfies

$$\int_{\partial K_n} \left( \frac{\partial}{\partial n_x} \left( \frac{1}{\omega_n(2-n)|x-y|^{n-2}} + \frac{1}{\omega_n(2-n)|y|^{n-2}|x-y^*|^{n-2}} + \alpha |x|^2 \right) \right) ds_x = 0,$$

where  $ds_x$  stands for the areal element of  $\partial K_n$ .

Since we have

$$\int_{\partial K_n} \frac{\partial}{\partial n_x} \left( \frac{1}{\omega_n(2-n)|x-y|^{n-2}} \right) ds_x = \begin{cases} 0 & y \notin \overline{K_n} \\ \frac{1}{2} & y \in \partial K_n \\ 1 & y \in K_n \end{cases}$$

and

$$\int_{\partial K_n} \frac{\partial}{\partial n_x} |x|^2 ds_x = 2\omega_n,$$

so the Neumann problem (2.7)(2.8) is solvable if and only if

$$\alpha = -\frac{1}{2\omega_n} \quad (2.10)$$

Thus the problem (2.1)(2.2) is solvable if and only if

$$c = -\frac{n}{\omega_n} \quad (2.11)$$

Now we look for the solutions of the problem (2.7)(2.8), where  $\alpha$  is defined by (2.10). Suppose that there exists a solution for the problem (2.7)(2.8) in the following form

$$h(x, y) = \sum_{k=0}^{\infty} a_k(y) |x|^k C_k^{\frac{n-2}{2}}(\cos \theta), \quad (2.12)$$

whereby  $C_k^{\frac{n-2}{2}}(t)$  are Gegenbauer polynomials,  $\theta$  denotes the angle between the radial directions  $ox$  and  $oy$ ,  $a_k(y)$  are functions of  $y$  to be determined. It is clear that the series (2.12) is harmonic in  $\Omega$  if it is uniformly convergent in  $\Omega$ . By the boundary condition (2.8) now we determine functions  $a_k(y)$ . Notice that, for  $|x| > |y|$ , we have

$$\frac{1}{|x-y|^{n-2}} = \sum_{k=0}^{\infty} C_k^{\frac{n-2}{2}}(\cos \theta) |y|^k |x|^{-(n+k-2)}$$

and

$$\frac{1}{|y|^{n-2} |x-y^*|^{n-2}} = \sum_{k=0}^{\infty} C_k^{\frac{n-2}{2}}(\cos \theta) |x|^k |y|^k.$$

On the one side, by (2.8) and (2.10) we have

$$\begin{aligned} \frac{\partial}{\partial n_x} h|_{\partial K_n} &= -\frac{1}{\omega_n(2-n)} \left( \sum_{k=0}^{\infty} (-(n+k-2)) C_k^{\frac{n-2}{2}}(\cos \theta) |y|^k \right. \\ &\quad \left. + \sum_{k=1}^{\infty} k C_k^{\frac{n-2}{2}}(\cos \theta) |y|^k \right) - 2\alpha \\ &= \sum_{k=1}^{\infty} \left( -\frac{1}{\omega_n} \right) C_k^{\frac{n-2}{2}}(\cos \theta) |y|^k. \end{aligned}$$

And on the other side, by (2.12) we have

$$\frac{\partial}{\partial n_x} h|_{\partial K_n} = \sum_{k=1}^{\infty} k a_k(y) C_k^{\frac{n-2}{2}}(\cos \theta).$$

Therefore we have

$$a_k(y) = -\frac{1}{k \omega_n} |y|^k, \quad k = 1, 2, \dots$$

Notice  $C_0^{\frac{n-2}{2}}(\cos \theta) = 1$ , so we have

$$h(x, y) = \sum_{k=1}^{\infty} \frac{-1}{k \cdot \omega_n} C_k^{\frac{n-2}{2}}(\cos \theta) |x|^k |y|^k + a_0(y). \quad (2.13)$$

It is easy to see that the series (2.13) is uniformly convergent in  $K_n$  with respect to  $x$  while  $y \in K_n$ . Now we can get the following proposition.

**Proposition 2.1** Let  $K_n$  be the unit ball in  $R^n (n \geq 3)$ . Then the function  $N_n(x, y)$

$$\begin{aligned} N_n(x, y) = & \frac{1}{(2-n)\omega_n |x-y|^{n-2}} + \frac{1}{(2-n)\omega_n |y|^{n-2} |x-y^*|^{n-2}} \\ & - \frac{1}{2\omega_n} (|x|^2 + |y|^2) - \frac{1}{\omega_n} \sum_{k=1}^{\infty} C_k^{\frac{n-2}{2}}(\cos \theta) \frac{|x|^k |y|^k}{k} \\ & - \frac{n^2 + n + 2}{\omega_n(4-n^2)} \end{aligned} \quad (2.14)$$

has the following properties:

$$i) \Delta_x N_n(x, y) = \delta(x-y) - \frac{n}{\omega_n}, \quad x \in K_n, \quad y \in K_n$$

$$ii) \frac{\partial}{\partial n_x} N_n(x, y) = 0, \quad x \in \partial K_n, \quad y \in K_n$$

$$iii) N_n(x, y) = N_n(y, x),$$

$$iv) \int_{K_n} N_n(x, y) dx = 0, \quad y \in K_n$$

whereby  $y^*$  is the symmetric point of  $y$  to the unit sphere,  $y^* = \frac{y}{|y|^2}$ ,  $dx$  is the volume element of  $K_n$ . The function  $N_n(x, y)$  is called the Neumann's function for Laplacian over the unit ball in  $R^n (n \geq 3)$ .

Indeed from above inference the properties (i)(ii)(iii) follow immediately.

Now we put, for  $y \in K_n$ ,

$$\begin{aligned} f_1(y) &= \int_{K_n} \frac{1}{(2-n)\omega_n |x-y|^{n-2}} dx, \\ f_2(y) &= \int_{K_n} \frac{1}{(2-n)\omega_n |y|^{n-2} |x-y^*|^{n-2}} dx, \\ f_3(y) &= \int_{K_n} \left( -\frac{1}{\omega_n} \sum_{k=1}^{\infty} C_k^{\frac{n-2}{2}}(\cos \theta) \frac{|x|^k |y|^k}{k} \right) dx. \end{aligned}$$

Notice that the function  $f_1(y)$  satisfies the Poisson equation  $\Delta f_1(y) = 1$ ,

$f_1(0) = \frac{1}{2(2-n)}$  and  $f_1(y)$  is only a function of  $|y|$ . Then we can easily get

$$f_1(y) = \frac{1}{2(2-n)} + \frac{1}{2n} |y|^2, \quad y \in K_n. \quad (2.15)$$

Notice also that  $f_1(y)$  is continuous in the whole space  $\mathbb{R}^n$ , so  $f_1(y) = \frac{1}{n(2-n)}$ , for  $|y|=1$ . Moreover,  $f_1(y)$  is harmonic in  $\mathbb{R}^n \setminus \overline{K_n}$  and vanishes at infinity. So we can also get

$$f_1(y) = \frac{1}{n(2-n)|y|^{n-2}}, \quad y \in \mathbb{R}^n \setminus \overline{K_n}. \quad (2.15')$$

Therefore we have

$$\begin{aligned} f_2(y) &= \frac{1}{|y|^{n-2}} \quad f_1(y^*) = \frac{1}{n(2-n)|y|^{n-2}|y^*|^{n-2}} \\ &= \frac{1}{n(2-n)}, \quad y \in K_n \end{aligned} \quad (2.16)$$

We also have

$$-\int_{K_n} \frac{1}{2\omega_n} (|x|^2 + |y|^2) dx = -\frac{1}{2(n+2)} - \frac{1}{2n} |y|^2. \quad (2.17)$$

Next we calculate  $f_3(y)$  for  $y \in K_n$ . Since the series  $h_3(y)$

$$h_3(y) = -\frac{1}{\omega_n} \sum_{k=1}^{\infty} C_k \frac{n-2}{2} (\cos \theta) \frac{|x|^k |y|^k}{k}$$

is harmonic in  $K_n$  with respect to  $y$  while  $x \in K_n$ , moreover,

$$\begin{aligned} \frac{\partial}{\partial n_y} h_3(y) \Big|_{y \in \partial K_n} &= -\frac{1}{\omega_n} \sum_{k=1}^{\infty} C_k \frac{n-2}{2} (\cos \theta) |x|^k \\ &= -\frac{1}{\omega_n} \left( \frac{1}{(1-2|x|\cos \theta + |x|^2)^{n-2/2}} - 1 \right). \end{aligned}$$

Then we have

$$\begin{aligned} \frac{\partial}{\partial n_y} f_3(y) \Big|_{y \in \partial K_n} &= -\frac{1}{\omega_n} \int_{K_n} \left( \frac{1}{(1-2|x|\cos \theta + |x|^2)^{n-2/2}} - 1 \right) dx \\ &= -(2-n)f_1(1) + \frac{1}{n} = 0, \end{aligned}$$

and  $f_3(y)$  is harmonic in  $K_n$ . Therefore we get

$$f_3(y) = f_3(o) = 0, \quad y \in K_n \quad (2.18)$$

From (2.15)-(2.18) the property (iv) follows immediately. Thus this proposition has been proved.

In addition, it is easy to see that the Neumann's function having the properties (i)-(iv) is unique. By above Neumann's function we can get the following proposition in a classical way.

Proposition 2.2 Let  $K_n$  be the unit ball in  $\mathbb{R}^n (n \geq 3)$ . Let  $f(x)$  be a continuous and differentiable function in  $\overline{K}_n$ . Then the Neumann problem for Poisson equation

$$\Delta u(x) = f(x), \quad x \in K_n \quad (2.19)$$

$$\frac{\partial}{\partial n} u(x) = 0, \quad x \in \partial K_n \quad (2.20)$$

is solvable in the classical sense if and only if  $f(x)$  satisfies

$$\int_{K_n} f(x) dx = 0 \quad (2.21)$$

and the solution may be represented in the following formula

$$u(x) = \int_{K_n} N_n(x, y) f(y) dy + c, \quad (2.22)$$

whereby the function  $N_n(x, y)$  is the Neumann's function described in the proposition 2.1 and  $c$  is an arbitrary real constant.

### 3. Boundary value problem of Neumann type

We first introduce the subalgebra  $H^{(i)}(R_{o, n-1})$  of  $R_{o, n-1}$ . Fixing  $i \in \{1, 2, \dots, n\}$  and considering for  $i > 2$  the subalgebra  $H^{(i)}(R_{o, n-1})$  of  $R_{o, n-1}$  is generated by the basis elements  $e_2, e_3, \dots, e_{i-1}$ , then  $H^{(i)}(R_{o, n-1})$  is isomorphic to the Clifford algebra  $R_{o, i-2}$  for  $R^{o, i-2}$ . For  $i=1, 2$ , we put  $H^{(i)}(R_{o, n-1}) = R$ .

Now define for  $i \in \{1, 2, \dots, n\}$  fixed the set  $I_i$  by

$$I_i = \begin{cases} \{i+1, \dots, n\}, & \text{if } i \in \{1, 2, \dots, n-1\} \\ \emptyset & \text{if } i=n. \end{cases}$$

And let us introduce the projective operator  $X^{(i)}$  of  $R_{0,n-1}$  onto its subspace  $H^{(i)}(R_{0,n-1}) e_{I_i}$ . Then we have for each  $a = \sum_A a_A e_A \in R_{0,n-1}$  that

$$X^{(i)} a = \sum_C a_C e_C$$

where  $C$  runs over all ordered subset of the form  $C = D \cup I_i$  with  $D \subset \{2, \dots, i-1\}$ , while for  $i=1$

$$X^{(1)}(a) = a_2 \dots e_2 \dots e_n.$$

Notice in particular that for  $i=n$

$$X^{(n)}(R_{0,n-1}) = H^{(n)}(R_{0,n-1}) e_1 = H^{(n)}(R_{0,n-1}),$$

i.e.  $X^{(n)}(R_{0,n-1})$  is the clifford subalgebra of  $R_{0,n-1}$  generated by the basis elements  $e_2, \dots, e_{n-1}$ , where

$$R_{0,n-1} = X^{(n)}(R_{0,n-1}) \oplus X^{(n)}(R_{0,n-1}) e_n. \quad (3.1)$$

Now consider the main involution  $a \rightarrow \hat{a}$  and the conjugation  $a \rightarrow \bar{a}$  on  $R_{0,n-1}$ , i.e. if  $a = \sum_A a_A e_A$ , then

$$\hat{a} = \sum_A a_A \hat{e}_A, \quad \bar{a} = \sum_A a_A \bar{e}_A,$$

whereby for each  $e_A = e_{\alpha_2} \dots e_{\alpha_h}$ ,

$$\hat{e}_A = (-1)^{\#A} e_A \quad \text{and} \quad \bar{e}_A = (-1)^{\#A} e_{\alpha_h} \dots e_{\alpha_2},$$

$\#A$  denoting the cardinality of  $A$ . Then for each  $d \in X^{(n)}(R_{0,n-1})$

$$d \cdot e_n = e_n \hat{d}.$$

We obtain by means of (3.1) that an arbitrary  $a \in R_{0,n-1}$  may be written as

$$a = b + c e_n = b + e_n \hat{c}, \quad b, c \in X^{(n)}(R_{0,n-1}).$$

Putting  $\hat{c} = Y^{(n)}(a)$ , we so have for all  $a \in R_{0,n-1}$  that

$$a = X^{(n)}(a) + e_n Y^{(n)}(a). \quad (3.2)$$

Notice that for  $n=2$ , we obtain that, since  $X^{(2)}(R_{0,1})$  is isomorphic to  $R$ , any  $a \in R_{0,1}$  admits the decomposition

$$a = b + e_2 c, \quad b, c \in R \quad (3.3)$$

whereby  $e_2^2 = -1$ . The relation (3.3) makes us remind of the well known result that  $R_{0,1}$  is isomorphic (as a real algebra) to  $C$ . But at the same time it illustrates how the decomposition obtained in (3.2) of an arbitrary element  $a$  in  $R_{0,n-1}$  is a generalization of the classical representation of a complex number  $a$  in the form  $a=x+iy$ ,  $x, y \in R$ .

Now let in a classical way  $R^i$  ( $i \geq 2$ ) be identified with subspace of  $R^n$  given by these elements  $x=(x_1, \dots, x_n) \in R^n$  for which  $x_{i+1} = \dots = x_n = 0$ . Then by means of the isomorphism between  $R^n$  and  $Re_1 \oplus R^{0,n-1}$ ,

$$x=(x_1, \dots, x_n), \quad z_x^{(n)} = \sum_{j=1}^n x_j e_j,$$

we may identify  $R^i$  with  $Re_1 \oplus R^{0,i-1}$ . An arbitrary element of  $R^i$  ( $i \geq 2$ ) may therefore be written as  $z_x^{(i)} = x_1 e_1 + x_2 e_2 + \dots + x_i e_i$ . Notice that for  $i=2, \dots, n$ ,

$$|z_x^{(i)}|^2 = \overline{z_x^{(i)}} z_x^{(i)} = z_x^{(i)} \overline{z_x^{(i)}} = \sum_{j=1}^i x_j^2.$$

Now we also consider the generalized Cauchy-Riemann operator  $\bar{\partial}_i$  and its conjugate operator  $\partial_i$  in  $R^i$ ,  $i=2, \dots, n$ , i.e.

$$\bar{\partial}_i = \sum_{j=1}^i e_j \frac{\partial}{\partial x_j} = e_1 \frac{\partial}{\partial x_1} + \sum_{k=2}^i e_k \frac{\partial}{\partial x_k}$$

and

$$\partial_i = \overline{\bar{\partial}_i} = e_1 \frac{\partial}{\partial x_1} - \sum_{k=2}^i e_k \frac{\partial}{\partial x_k}.$$

Let  $K_n^{(i)}$  and  $\partial K_n^{(i)}$  be the unit ball and unit sphere in  $R^i$  respectively, i.e.

$$K_n^{(i)} = \{x=(x_1, x_2, \dots, x_n) : \sum_{j=1}^i x_j^2 < 1 \text{ and } \sum_{j=i+1}^n x_j^2 = 0\}$$

and

$$\partial K_n^{(i)} = \{x=(x_1, x_2, \dots, x_n) : \sum_{j=1}^i x_j^2 = 1 \text{ and } \sum_{j=i+1}^n x_j^2 = 0\}.$$

Now we consider the following integral operator  $T_i$  which are also called the Pompeiu-operator and may be defined on the space of  $R_{0,i-1}$ -valued functions  $g$

$$(T_i g)(z_x^{(i)}) = -\frac{1}{\omega_i} \int_{K_n^{(i)}} \frac{z_x^{(i)} - z_y^{(i)}}{|z_x^{(i)} - z_y^{(i)}|^i} g(z_y^{(i)}) dy_1 \dots dy_i, \quad (3.4)$$

$i=2,3,\dots,n-1,$

whereby  $\omega_i$  is the area of  $\partial K_n^{(i)}$ , and  $g$  is of the class  $C_\alpha$  ( $0 < \alpha < 1$ ) in  $\overline{K_n^{(i)}}$ .

By [4] it is well known that

$$\partial_i (T_i g) = g. \quad (3.5)$$

Now let us formulate the following basic lemma.

Lemma 3.1 The nonhomogeneous generalized Cauchy-Riemann equation

$$\overline{\partial}_n w = f, \quad \text{in } K_n^{(n)} \quad (3.6)$$

is equivalent to the following equations

$$\overline{\partial}_{n-1} X^{(n)}_w - \frac{\partial}{\partial x_n} Y^{(n)}_w = X^{(n)}_f, \quad \text{in } K_n^{(n)} \quad (3.7)$$

$$\frac{\partial}{\partial x_n} X^{(n)}_w + \partial_{n-1} Y^{(n)}_w = Y^{(n)}_f, \quad \text{in } K_n^{(n)} \quad (3.8)$$

whereby  $(X^{(n)}_w)(x) = X^{(n)}(w(x))$  and  $(Y^{(n)}_w)(x) = Y^{(n)}(w(x))$ .

Proof As  $\overline{\partial}_{n-1}(e_n Y^{(n)}_w) = e_n \partial_{n-1}(Y^{(n)}_w)$ , we have

$$\begin{aligned} \overline{\partial}_n w &= \left( \overline{\partial}_{n-1} + e_n \frac{\partial}{\partial x_n} \right) (X^{(n)}_w + e_n Y^{(n)}_w) \\ &= \left( \overline{\partial}_{n-1} X^{(n)}_w - \frac{\partial}{\partial x_n} Y^{(n)}_w \right) + e_n \left( \frac{\partial}{\partial x_n} X^{(n)}_w + \partial_{n-1} Y^{(n)}_w \right) \\ &= X^{(n)}_f + e_n Y^{(n)}_f. \end{aligned}$$

Thus this lemma is proved.

Theorem 3.2 Let  $f(x)$  be a continuous differentiable  $R_{0,n-1}$ -valued function in  $\overline{K_n^{(n)}}$ . Then the boundary value problem of Neumann type

$$\overline{\partial}_n w = f, \quad x \in K_n^{(n)} \quad (3.9)$$

$$X^{(n)}(\frac{\partial}{\partial n} w) = 0, \quad x \in \partial K_n^{(n)} \quad (3.10)$$

is solvable if and only if  $f(x)$  satisfies the following integral-differential relation

$$\int_{K_n^{(n)}} (\partial_{n-1}(X^{(n)}f) + \frac{\partial}{\partial x_n} (Y^{(n)}f)) dx = 0. \quad (3.11)$$

And the solution may be represented in the following form

$$\begin{aligned} w &= X^{(n)}w + e_n Y^{(n)}w, \\ X^{(n)}w &= \int_{K_n^{(n)}} N_n(x, y) (\partial_{n-1}(X^{(n)}f) + \frac{\partial}{\partial y_n} (Y^{(n)}f))(y) dy + c, \\ Y^{(n)}w &= \int_0^{x_n} (\overline{\partial}_{n-1} X^{(n)}w - X^{(n)}f) dx_n \\ &\quad + T_{n-1}((-\frac{\partial}{\partial x_n} X^{(n)}w + Y^{(n)}f) |_{x_n=0}) + \tilde{h}(z_x^{(n-1)}), \end{aligned} \quad (3.12) \quad (3.13)$$

where  $c$  is an arbitrary  $X^{(n)}(\mathbb{R}_{0,n-1})$ -valued constant and  $\tilde{h}(z_x^{(n-1)})$  is an arbitrary  $X^{(n)}(\mathbb{R}_{0,n-1})$ -valued function satisfying  $\partial_{n-1}\tilde{h}(z_x^{(n-1)})=0$ .

**Proof** If there exists a solution  $w(x)$  for the problem (3.9)(3.10) in the classical sense, then  $w(x)$  should satisfies the following problem

$$\partial_n(\overline{\partial}_n w) = \partial_n f, \quad x \in K_n^{(n)} \quad (3.14)$$

$$\frac{\partial}{\partial n} (X^{(n)}w) = 0, \quad x \in \partial K_n^{(n)} \quad (3.15)$$

Then we have that  $X^{(n)}w$  should satisfies the following problem

$$\Delta_n X^{(n)}w = \partial_{n-1}(X^{(n)}f) + \frac{\partial}{\partial x_n} (Y^{(n)}f), \quad x \in K_n^{(n)} \quad (3.16)$$

$$\frac{\partial}{\partial n} (X^{(n)}w) = 0, \quad x \in \partial K_n^{(n)}. \quad (3.17)$$

By proposition 2.2 then we get (3.11)(3.12). By lemma 3.1 and (3.7) we have

$$Y^{(n)}w = \int_0^{x_n} (\overline{\partial}_{n-1} X^{(n)}w - X^{(n)}f) dx_n + h(z_x^{(n-1)}),$$

where  $h(z_x^{(n-1)})$  is an arbitrary  $X^{(n)}(R_{0,n-1})$ -valued function depending only on  $x_1, \dots, x_{n-1}$ . Substituting into (3.8), we get

$$\frac{\partial}{\partial x_n} X^{(n)}_w + \int_0^{x_n} (\Delta_{n-1} X^{(n)}_w - \partial_{n-1}(X^{(n)}_f)) dx_n + \partial_{n-1} h(z_x^{(n-1)}) = Y^{(n)}_f \quad .$$

By (3.16) we have

$$\Delta_{n-1} X^{(n)}_w = - \frac{\partial^2}{\partial x_n^2} X^{(n)}_w + \partial_{n-1}(X^{(n)}_f) + \frac{\partial}{\partial x_n} (Y^{(n)}_f),$$

hence we obtain

$$\frac{\partial}{\partial x_n} X^{(n)}_w \Big|_{x_n=0} - Y^{(n)}_f \Big|_{x_n=0} + \partial_{n-1} h(z_x^{(n-1)}) = 0.$$

By (3.5) then we get

$$h(z_x^{(n-1)}) = T_{n-1}((- \frac{\partial}{\partial x_n} X^{(n)}_{w+Y^{(n)}_f} \Big|_{x_n=0}) + \tilde{h}(z_x^{(n-1)}),$$

whereby  $\tilde{h}(z_x^{(n-1)})$  is an arbitrary  $X^{(n)}(R_{0,n-1})$ -valued function satisfying the equation  $\partial_{n-1} \tilde{h}(z_x^{(n-1)}) = 0$ . Thus we get the formula (3.13).

Conversely if the function  $w(x)$  is defined by (3.12)(3.13), then we get immediately that the functions  $X^{(n)}_w$  and  $Y^{(n)}_w$  satisfy (3.7). By (3.12) we have that the function  $X^{(n)}_w$  satisfies (3.16), then we get

$$\begin{aligned} & \frac{\partial}{\partial x_n} X^{(n)}_w + \partial_{n-1} Y^{(n)}_w \\ &= \frac{\partial}{\partial x_n} X^{(n)}_w + \int_0^{x_n} \partial_{n-1} (- \frac{\partial}{\partial x_n} X^{(n)}_w - X^{(n)}_f) dx_n + (- \frac{\partial}{\partial x_n} X^{(n)}_{w+Y^{(n)}_f}) \Big|_{x_n=0} \\ &= \frac{\partial}{\partial x_n} X^{(n)}_w + \int_0^{x_n} (- \frac{\partial^2}{\partial x_n^2} X^{(n)}_w + \frac{\partial}{\partial x_n} (Y^{(n)}_f)) dx_n + (- \frac{\partial}{\partial x_n} X^{(n)}_{w+Y^{(n)}_f}) \Big|_{x_n=0} \\ &= Y^{(n)}_f \quad . \end{aligned}$$

Hence the functions  $X^{(n)}_w$  and  $Y^{(n)}_w$  also satisfy (3.8). By lemma 3.1 the function  $w$  defined by (3.12)(3.13) satisfies (3.9). Obviously (3.10) yields. Thus this theorem has been proved.

Now we introduce the integral-differential operator  $K_n$  defined on the space of continuous and differentiable  $R_{0,n-1}$ -valued function  $f(x)$

$$(\mathbb{K}_n f)(x) = (\mathbb{K}_n^{(1)} f)(x) + e_n (\mathbb{K}_n^{(2)} f)(x)$$

whereby

$$\begin{aligned} (\mathbb{K}_n^{(1)} f)(x) &= \int_{\mathbb{K}_n^{(n)}} N_n(x, y) \left( \partial_{n-1} (X^{(n)} f) + \frac{\partial}{\partial y_n} (Y^{(n)} f) \right)(y) dy, \\ (\mathbb{K}_n^{(2)} f)(x) &= \int_0^{x_n} \left( \bar{\partial}_{n-1} (\mathbb{K}_n^{(1)} f) - X^{(n)} f \right) dx + T_{n-1} \left( \left( -\frac{\partial}{\partial x_n} (\mathbb{K}_n^{(1)} f) + Y^{(n)} f \right) \Big|_{x_n=0} \right). \end{aligned}$$

Then we can get

Corollary 3.3 Under the assumptions of theorem 3.2 the solution of the boundary value problem (3.9)(3.10) may be represented in the following form

$$w(x) = (\mathbb{K}_n f)(x) + c + h_{n-1}(z_x^{(n-1)}) e_n,$$

where  $c$  is an arbitrary  $X^{(n)}(\mathbb{R}_{0,n-1})$ -valued constant and  $h_{n-1}(z_x^{(n-1)})$  is an arbitrary  $X^{(n)}(\mathbb{R}_{0,n-1})$ -valued function satisfying  $\bar{\partial}_{n-1} h_{n-1}(z_x^{(n-1)}) = 0$ .

Remark Making use of the result in [11], the solution of the problem (3.9)(3.10) can be determined uniquely on condition that  $(X^{(n)} w)(0)$ , the functions  $X^{(i)} w$  on  $\partial \mathbb{K}_n^{(i)}$  ( $i = n-1, \dots, 2$ ) and  $(X^{(1)} w)(0)$  are also given.

#### REFERENCES

1. Brackx F., Delanghe R. and Sommen F. Clifford Analysis, Pitman, London, 1982.
2. Chisholm J.S.R., Commen A.K., Eds. Clifford Algebras and Their Applications in Mathematical Physics, Reidel, Dordrecht, 1986.
3. Folland G.B. Introduction to Partial Differential Equations, Princeton, New Jersey, 1976.
4. Gilbert R.P., Buchanan J.L. First Order Elliptic Systems: A Function Theoretic Approach, Academic press, New York, 1983.
5. Hestenes D., Sobczyk G. Clifford Algebra to Geometric Calculus: A Unified Language for Mathematics and Physics, Reidel, Dordrecht, 1984.
6. Iftimie V. Functions hypercomplexes, Bull. Soc. sci. Math. R.S. Roumanie 9(1965)279-332.

7. Porteous I.R. Topological Geometry , Van Nostrand, London, 1969.
8. Stern I. Boundary value problems for generalized Cauchy-Riemann system in the space , A lecture at "5<sup>th</sup> conference on complex analysis and its application" in Halle,DDR.
9. Xu Zhenyuan On Riemann boundary value problems for regular function with values in a Clifford algebra , Kexue Tongbao (Science Bulletin) (18)32(1987),1294-1295.
10. Xu Zhenyuan, Chen Jin. On Riemann-Hilbert problem for regular function with values in a Clifford algebra , Kexue Tongbao (Science Bulletin) (10)33(1988),874-876.
11. Xu Zhenyuan, Chen Jin, Zhang Wanguo AHarmonic conjugate of the Poisson kernel and a boundary value problem for monogenicfunctions in the unit ball of  $R^n(n \geq 2)$ , to appear in Simon Stevin.

DEPARTMENT OF MATHEMATICS

FUDAN UNIVERSITY

SHANGHAI, PEOPLE'S REPUBLIC OF CHINA