

Manoj Dhake; Sachin Ballal; Vilas S. Kharat; Rupesh S. Shewale
On forbidden configuration of pseudomodular lattices

Mathematica Bohemica, Vol. 150 (2025), No. 1, 99–108

Persistent URL: <http://dml.cz/dmlcz/152878>

Terms of use:

© Institute of Mathematics CAS, 2025

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON FORBIDDEN CONFIGURATION OF PSEUDOMODULAR LATTICES

MANOJ DHAKE, Pune, SACHIN BALLAL, Hyderabad,
VILAS KHARAT, RUPESH S. SHEWALE, Pune

Received August 17, 2023. Published online July 1, 2024.

Communicated by Simion Breaz

Abstract. We characterize the pseudomodular lattices by means of a forbidden configuration.

Keywords: forbidden configuration; pseudomodular lattice; semimodular lattice

MSC 2020: 06C10, 06C99

1. INTRODUCTION AND PRELIMINARIES

Dress and Lovász in [4] studied full algebraic matroids of finite ranks for modularity of the flats. They characterized the existence of flats using the rank function and quasi-intersection. Björner and Lovász in [2] introduced a class of pseudomodular lattices as a generalization of modular lattices to contain full algebraic combinatorial geometries; see also [3]. A semimodular lattice L of finite length is said to be pseudomodular if every pair of elements of L has a pseudointersection. The class of pseudomodular lattices forms a subclass of the class of semimodular lattices and contains all modular lattices of finite length. Characterizations of classess of lattices by means of the non-existence of certain sublattices called forbidden configurations are available in the literature, such as the classess of distributive lattices, modular lattices, semimodular lattices, etc. In this paper, we establish a characterization by means of a forbidden sublattice for the class of pseudomodular lattices.

We give here some definitions and notations for ready reference; see Birkhoff [1], Grätzer [5], Haskins and Gudder [6], Stern [7], etc.

This work is supported by University Grants Commission, under JRF, Govt. of India.

Let P be a nonempty poset and $x, y \in P$. If $x \leq y$, then the length of an interval $[x, y]$, denoted by $lt[x, y]$, is the supremum of the lengths of the chains in $[x, y]$. The *height* or *rank* $r(x)$ of an element x of a poset P bounded below is the length of the interval $[0, x]$.

A lattice L is (*upper*) *semimodular* if $a \wedge b$ is a lower cover of a . Then b is a lower cover of $a \vee b$, for $a, b \in L$. A lattice L is said to be modular if the following condition (M) holds.

(M): $c \vee (a \wedge b) = (c \wedge a) \vee b$ for all $a, b, c \in L$ with $c \leq b$.

Definition 1.1 ([1]). The *graded poset* is defined as a poset P with a function $g: P \rightarrow \mathbb{Z}$ such that:

- (i) $x > y$ implies $g[x] > g[y]$ and
- (ii) if $x \prec y$, then $g[x] = g[y] + 1$.

Note that any semimodular lattice of finite length is graded by its rank function. Following is the definition due to Björner and Lovász [2], see also [3].

Definition 1.2 ([2]). Let L be a semimodular lattice of finite length, and denote by $r(x)$ the rank function (height function) of L . For each $x, y \in L$ let $P_{x,y} = \{z \leq y: r(x \vee z) - r(z) = r(x \vee y) - r(y)\}$. If the set $P_{x,y}$ has a unique least element, then we call this the *pseudointersection* of x and y and denote it by $x \downarrow y$.

A semimodular lattice of finite length is called *pseudomodular* if every pair of its elements has the pseudointersection.

Remark 1.3 ([2]). The set $P_{x,y}$ lies in the interval $[x \wedge y, y]$ and is dual order ideal in $[x \wedge y, y]$.

Lemma 1.4 ([2]). For any two elements x and y in a semimodular lattice L , the following are equivalent:

- (i) x and y form a modular pair, i.e., $r(x \vee y) + r(x \wedge y) = r(x) + r(y)$.
- (ii) $x \downarrow y$ exists and $x \downarrow y \leq x$.
- (iii) $x \downarrow y$ exists and $x \downarrow y = x$.
- (iv) $x \wedge y \in P_{x,y}$.

Lemma 1.5 ([2]). For any two elements x and y of a semimodular lattice L , the following are equivalent:

- (i) $x \downarrow y$ exists, i.e., $P_{x,y}$ has a unique least element.
- (ii) $P_{x,y}$ is closed under meets.
- (iii) If $u, v, z \in P_{x,y}$ and z covers u and v , then $u \wedge v \in P_{x,y}$.

A subset I of a poset P is an order ideal if $x \in I$ and $y \leq x$ imply $y \in I$.

2. MAIN RESULT

We have a forbidden characterization of pseudomodular lattices in the following theorem. In what follows, a sublattice S of a lattice L is said to be *cover-preserving* in L if for $a, b \in S$, $a \prec b$ in S implies $a \prec b$ in L , see [7].

Theorem 2.1. *Let L be a semimodular lattice of finite length. Then L is pseudomodular if and only if L does not contain a cover preserving sublattice isomorphic to the lattice as depicted in Figure 1.*




Figure 1.

Proof of Theorem 2.1. Let L be a semimodular lattice of finite length. If L contains a sublattice isomorphic to the lattice as depicted in Figure 1, then the elements c_4, b_4, b_6 belong to P_{b_1, c_4} . Thus $r(b_1 \vee c_4) - r(c_4) = r(b_1 \vee b_4) - r(b_4) = r(b_1 \vee b_6) - r(b_6)$, but $r(b_1 \vee (b_4 \wedge b_6)) - r(b_4 \wedge b_6) \neq r(b_1 \vee b_6) - r(b_6)$. Therefore $b_4 \wedge b_6 \notin P_{b_1, c_4}$, which implies that P_{b_1, c_4} does not have the least element and so the pseudointersection of b_1 and c_4 does not exist. Therefore, L is not a pseudomodular lattice.

Conversely, suppose that L is a semimodular lattice of finite length which is not pseudomodular. Then there exists a pair of elements $x, y \in L$ such that $P_{x, y} = \{z \leq y : r(x \vee z) - r(z) = r(x \vee y) - r(y)\}$ does not have the least element. Or equivalently, we have a pair x, y in L whose meet does not belong to $P_{x, y}$.

Consider a pair z_1, z_2 in $P_{x, y}$ with minimal height whose meet does not belong to $P_{x, y}$. Since $y, z_1, z_2 \in P_{x, y}$, we have $r(x \vee y) - r(y) = r(x \vee z_1) - r(z_1) = r(x \vee z_2) - r(z_2)$. Without loss of generality we assume that $lt[x \wedge y, x \vee y]$ is minimum, i.e., for $u, v \in L$, if $lt[u \wedge v, u \vee v] < lt[x \wedge y, x \vee y]$, then $P_{u, v}$ has the least element. We also assume that $x, y \in L$ is a pair such that for $x \wedge y < u < x$, the set $P_{u, y}$ has the least element.

If $x \leq y$, then x, y becomes a modular pair and by Lemma 1.4, $x \wedge y \in P_{x,y}$, which is nothing but the pseudointersection of x and y , a contradiction to the assumption.

Similarly, if $x > y$, then also by Lemma 1.4, we have a contradiction. Consequently, we must have $x \parallel y$.

If $x \wedge y \prec x$, then by semimodularity we have $y \prec x \vee y$ and thus $r(x) - r(x \wedge y) = 1 = r(x \vee y) - r(y)$ and $P_{x,y} = \{z \leq y: r(x \vee z) - r(z) = r(x \vee y) - r(y)\} = \{z \leq y: r(x \vee z) - r(z) = 1\}$. Also, as $x \wedge y \leq y$ and $r(x \vee (x \wedge y)) - r(x \wedge y) = r(x) - r(x \wedge y) = 1$, we have $x \wedge y \in P_{x,y}$ and so $x \wedge y$ is the least element of $P_{x,y}$, a contradiction, and therefore we must have $x \wedge y \not\prec x$.

Consider an element q such that $x \wedge y < q < x$ and without loss of generality, we consider $x \wedge y < q \prec x$. If $x \wedge y \prec z_1$, then by semimodularity we have $q \prec q \vee z_1$ and also $x \prec x \vee z_1$. Therefore, $r(x \vee (x \wedge y)) - r(x \wedge y) = r(x \vee z_1) - r(z_1)$ and consequently, $x \wedge y \in P_{x,y}$, a contradiction, and so, we must have $x \wedge y \not\prec z_1$. Similarly, we have to have $x \wedge y \not\prec z_2$.

Now, since $x \wedge y \not\prec z_1$ and $x \wedge y \not\prec z_2$, there exist q_1 and q_2 such that $z_1 \wedge z_2 < q_1 < z_1$ and $z_1 \wedge z_2 < q_2 < z_2$ and without loss of generality, we consider $x \wedge y < q_1 \prec z_1$ and $x \wedge y < q_2 \prec z_2$.

Consider the set $\{x, y, x \vee y, x \wedge y, z_1, z_2, x \vee z_1, x \vee z_2, q, q_1, q_2, q \vee q_1, q \vee q_2, q_1 \vee q_2, q \vee q_1 \vee q_2\}$ and we contend that these elements are distinct and also the set forms a cover preserving sublattice of L . Note that by the choice $x, y, x \vee y, x \wedge y, z_1, z_2, x \vee z_1, x \vee z_2, q, q_1$ and q_2 are distinct elements. For the other elements, we have the following.

Claim 2.2. $(x \vee z_1) \wedge (x \vee z_2) = x$.

Proof. Suppose that $(x \vee z_1) \wedge (x \vee z_2) > x$. As $q_1 \prec z_1$, by semimodularity we have $q_1 \vee x \prec z_1 \vee x$. If $q_1 \vee x \prec z_1 \vee x$, then $r(q_1 \vee x) - r(q_1) = r(z_1 \vee x) - r(z_1) = r(x \vee y) - r(y) = r(x \vee z_2) - r(z_2)$, which implies $q_1 \in P_{x,y}$, a contradiction, and so we must have $(x \vee z_1) \wedge (x \vee z_2) = x$. $\square$

Claim 2.3. $x \wedge y = z_1 \wedge z_2$.

Proof. Suppose $x \wedge y < z_1 \wedge z_2$. If $x_1 = x \vee (z_1 \wedge z_2)$, then we have $x_1 \leq (x \vee z_1) \wedge (x \vee z_2)$, which gives $x \vee y = x_1 \vee y$, $x \vee z_1 = x_1 \vee z_1$ and $x \vee z_2 = x_1 \vee z_2$. It follows that $r(x_1 \vee y) - r(y) = r(x_1 \vee z_1) - r(z_1) = r(x_1 \vee z_2) - r(z_2)$ and so $y, z_1, z_2 \in P_{x_1,y}$. Since $lt[x_1 \wedge y, x_1 \vee y] < lt[x \wedge y, x \vee y]$, we have $z_1 \wedge z_2 \in P_{x_1,y}$. Thus, $r(x_1 \vee (z_1 \wedge z_2)) - r(z_1 \wedge z_2) = r(x_1 \vee y) - r(y) = r(x \vee y) - r(y)$. Moreover, $z_1 \wedge z_2 \in P_{x,y}$, a contradiction, and so we must have $x \wedge y = z_1 \wedge z_2$. $\square$

Claim 2.4. $z_1 \vee z_2 = y$.

Proof. Suppose that $z_1 \vee z_2 < y$. If $y_1 = z_1 \vee z_2$, then $z_1 < y_1 < y$. Since $P_{x,y}$ is a dual order ideal, we have $y_1 \in P_{x,y}$ and so $r(x \vee y_1) - r(y_1) = r(x \vee y) - r(y)$, which gives $x \vee y \neq x \vee y_1$. Now consider the interval $[x \wedge y_1, x \vee y_1]$. Since $lt[x \wedge y_1, x \vee y_1] < lt[x \wedge y, x \vee y]$ and $z_1, z_2, y_1 \in P_{x,y_1}$, we have $z_1 \wedge z_2 \in P_{x,y_1}$. Thus $r(x \vee (z_1 \wedge z_2)) - r(z_1 \wedge z_2) = r(x \vee y_1) - r(y_1) = r(x \vee y) - r(y)$, which implies $z_1 \wedge z_2 \in P_{x,y}$, a contradiction, and so $z_1 \vee z_2 = y$. $\square$

Claim 2.5. $(x \vee z_1) \vee (x \vee z_2) = x \vee y$.

Proof. Observe that $(x \vee z_1) \vee (x \vee z_2) = x \vee (z_1 \vee z_2) = x \vee y$. $\square$

Claim 2.6. $q \vee y = x \vee y$.

Proof. Suppose that $q \vee y < x \vee y$. As $q \prec x$, by semimodularity we have $q \vee y \prec x \vee y$. In this case, $q \vee z_1 \prec x \vee z_1$ and $q \vee z_2 \prec x \vee z_2$. If $q \vee z_1 = x \vee z_1$ or $q \vee z_2 = x \vee z_2$, then this implies that $q \vee z_1 \vee y = x \vee z_1 \vee y = q \vee y = x \vee y$, which is not possible and so, $r(x \vee y) - r(y) = r(x \vee z_1) - r(z_1) = r(x \vee z_2) - r(z_2)$. Also, we have $r(q \vee y) - r(y) = r(q \vee z_1) - r(z_1) = r(q \vee z_2) - r(z_2)$, which implies $y, z_1, z_2 \in P_{q,y}$. Since $lt[q \wedge y, q \vee y] < lt[x \wedge y, x \vee y]$, we have $z_1 \wedge z_2 \in P_{q,y}$ and so, $r(q \vee (z_1 \wedge z_2)) - r(z_1 \wedge z_2) = r(q \vee y) - r(y) = r(x \vee y) - r(y) - 1$. Consequently, we have $r(q \vee (z_1 \wedge z_2)) + 1 - r(z_1 \wedge z_2) = r(x \vee y) - r(y)$. Now, since $q \prec x$ and $q \vee (z_1 \wedge z_2) = q$, we have $r(q \vee (z_1 \wedge z_2)) + 1 = r(x \vee (z_1 \wedge z_2))$, which gives $r(x \vee (z_1 \wedge z_2)) - r(z_1 \wedge z_2) = r(x \vee y) - r(y)$. This implies that $z_1 \wedge z_2 \in P_{x,y}$, a contradiction, and so we must have $y \vee q = x \vee y$. $\square$

Claim 2.7. $q \vee z_1 = x \vee z_1$.

Proof. Suppose that $q \vee z_1 < x \vee z_1$ and consider a chain of length n in $[z_1, x \vee z_1]$: $z_1 \prec p_1 \prec p_2 \prec \dots \prec p_{n-1} = q \vee z_1 \prec x \vee z_1$. By semimodularity, we have a chain in $[y, x \vee y]$: $y = z_1 \vee y \prec p_1 \vee y \preceq p_2 \vee y \preceq \dots \preceq p_n = q \vee z_1 \vee y = x \vee z_1 \vee y$ which is of length at most n , a contradiction to the fact that $r(q \vee y) - r(y) = r(q \vee z_1) - r(z_1)$. So we must have $q \vee z_1 = x \vee z_1$. $\square$

Claim 2.8. $q \vee z_2 = x \vee z_2$.

Proof. Is similar to that of Claim 2.7. $\square$

Claim 2.9. $x \vee q_1 = x \vee z_1$.

Proof. Suppose that $x \vee q_1 < x \vee z_1$. As $q_1 \prec z_1$, by semimodularity we have $x \vee q_1 \prec x \vee z_1$. This gives $r(x \vee z_1) - r(z_1) = r(x \vee q_1) - r(q_1)$. Thus, $q_1 \in P_{x,y}$, a contradiction, and so we must have $x \vee q_1 = x \vee z_1$. $\square$

Claim 2.10. $x \vee q_2 = x \vee z_2$.

Proof. Is similar to that of Claim 2.9. $\square$

Claim 2.11. $q_1 \vee z_2 = y$.

Proof. Suppose that $q_1 \vee z_2 < y$. As $q_1 \prec z_1$, by semimodularity we have $q_1 \vee z_2 \prec y$. If $y_1 = q_1 \vee z_2$, then $z_2 < y_1 < y$. Since $P_{x,y}$ is a filter, we have $y_1 \in P_{x,y}$, which implies $r(x \vee y_1) - r(y_1) = r(x \vee y) - r(y)$. Also, we have $q_1 \leq y_1 \leq x \vee y_1$ and $x \leq x \vee y_1$, therefore $x \vee q_1 \leq x \vee y_1$. Since $x \vee q_1 = x \vee z_1$, we have $z_1 \leq x \vee z_1 \leq x \vee y_1$. Also $z_2 \leq y_1 \leq x \vee y_1$ and therefore $z_1 \vee z_2 \leq x \vee y_1$. However, we have $z_1 \vee z_2 = y < x \vee y_1$, a contradiction, and so we must have $q_1 \vee z_2 = y$. $\square$

Claim 2.12. $q_2 \vee z_2 = y$.

Proof. Is similar to that of Claim 2.11. $\square$




Figure 2.

Claim 2.13. $q \vee q_1 < x \vee z_1$.

Proof. Suppose that $q \vee q_1 = x \vee z_1$. Note that $x \wedge y \not\leq q$ and $x \wedge y \not\leq q_1$; otherwise, $q_1 \prec q \vee q_1$, which is not true since $q_1 \prec z_1 < x \vee z_1$. Therefore there exists p_1 such that $x \wedge y \prec p_1 < q$. We have $p_1 \parallel z_1$, $p_1 \parallel q_1$ and $p_1 \vee q_1 \parallel z_1$. Now, if $(p_1 \vee q_1) \vee z_1 = x \vee z_1$, then by semimodularity we have $p_1 \vee q_1 \prec x \vee z_1$ and $z_1 \prec x \vee z_1$ and so $r(x \vee z_1) - r(z_1) = 1$. Also, as $q \vee y = x \vee y$, $q \vee z_1 = x \vee z_1$, $q \vee z_2 = x \vee z_2$ and $q < x$ and so, by assumption, $P_{q,y}$ must have the least element. We have $y, z_1, z_2 \in P_{q,y}$, $z_1 \wedge z_2 \in P_{q,y}$ but $r(q \vee (x \wedge y)) - r(x \wedge y) \neq 1$, a contradiction, and so we must have $q \vee q_1 < x \vee z_1$. $\square$

Claim 2.14. $q \vee q_2 < x \vee z_2$.

Proof. Is similar to that of Claim 2.13. $\square$

Claim 2.15. $x \wedge y \prec q$.

Proof. Suppose there exists an element p such that $x \wedge y \prec p < q$. It follows that $p \vee q_1 \leq q \vee q_1$. If $p \vee q_1 = q \vee q_1$, then by semimodularity we have $q_1 \prec q \vee q_1$, which is not true. Therefore $p \vee q_1 < q \vee q_1$ and by semimodularity we have $q_1 \prec p \vee q_1$ and similarly, $q_2 \prec p \vee q_2$. In this case, $(p \vee q_1) \vee z_1 \leq x \vee z_1$. We consider the following subcases:

(i) Suppose $(p \vee q_1) \vee z_1 = x \vee z_1$. By semimodularity we have $z_1 \prec x \vee z_1$ and $y \prec x \vee y$ and therefore $r(x \vee z_1) - r(z_1) = r(x \vee y) - r(y) = 1$. Since $r(x \vee y) - r(y) = r(x \vee z_2) - r(z_2)$, we have $r(x \vee z_2) - r(z_2) = 1$ and hence $z \prec x \vee z_2$. We also have $q \vee y = x \vee y$, $q \vee z_1 = x \vee z_1$, $q \vee z_2 = x \vee z_2$. Thus $y_1, z_1, z_2 \in P_{q,y}$ and by assumption, $P_{q,y}$ must have the least element, which gives $z_1 \wedge z_2 \in P_{q,y}$. Therefore $r(q \vee y) - r(y) = r(q \vee (z_1 \wedge z_2)) - r(z_1 \wedge z_2) = 1$, a contradiction to the fact that $r(q) - r(x \wedge y) > 1$, and therefore $(p \vee q_1) \vee z_1 \neq x \vee z_1$. Similarly, $(p \vee q_2) \vee z_2 \neq x \vee z_2$.

(ii) Suppose $(p \vee q_1) \vee z_1 < x \vee z_1$. Let $p_1 = (p \vee q_1) \vee z_1$. By semimodularity we have $z_1 \prec p_1$. If $p_1 \vee y = x \vee y$, then $y \prec x \vee y$, a contradiction to the fact that $r(x \vee y) - r(y) = r(x \vee z_1) - r(z_1)$, and therefore $p_1 \vee y < x \vee y$. By semimodularity we have $y \prec p_1 \vee y$. Similarly, for $p_2 = (p \vee q_2) \vee z_2$, we have $y \prec p_2 \vee y$.

In this case, $p_1 \vee y = p_2 \vee y = (p \vee q_2 \vee z_2) \vee y = (p \vee z_2) \vee y = p \vee y$. Let $y_1 = p_1 \vee y = p_2 \vee y = p \vee y$. Then $x \vee y_1 = x \vee y$, $x \wedge y < x \wedge y_1$ and $x \wedge y_1 \geq p$. Therefore $lt[x \wedge y_1, x \vee y_1] < lt[x \wedge y, x \vee y]$. As $r(x \vee y_1) - r(y_1) = r(x \vee p_1) - r(p_1) = r(x \vee z_1) - r(p_1) = r(x \vee p_2) - r(p_2) = r(x \vee z_2) - r(p_2)$, we have $y_1, p_1, p_2 \in P_{x,y_1}$. Hence P_{x,y_1} has the least element, which gives $p_1 \wedge p_2 \in P_{x,y_1}$. Thus $r(x \vee y_1) - r(y_1) = r(x \vee (p_1 \wedge p_2)) - r(p_1 \wedge p_2)$ and we have $p_1 \wedge p_2 \geq p$. Also, $p_1 \wedge p_2 \leq x \vee z_1$ and $p_1 \wedge p_2 \leq x \vee z_2$, which gives $p_1 \wedge p_2 \leq (x \vee z_1) \wedge (x \vee z_2)$, and so $p_1 \wedge p_2 \leq x$. In this case, $q \vee y = q \vee y_1 = x \vee y$, $q \vee z_1 = q \vee p_1 = x \vee z_1$, $q \vee z_2 = q \vee p_2 = x \vee z_2$ and $q \wedge y_1 > x \wedge y$, and so $y_1, p_1, p_2 \in P_{q,y_1}$. By assumption, $P_{q,y}$ has the least element and so $p_1 \wedge p_2 \in P_{q,y}$. Thus $r(q \vee (p_1 \wedge p_2)) - r(p_1 \wedge p_2) = r(x \vee z_1) - r(p_1)$. Since $(x \vee (p_1 \wedge p_2)) - r(p_1 \wedge p_2) = r(x \vee z_1) - r(p_1)$, which implies that $r(q \vee (p_1 \wedge p_2)) - r(p_1 \wedge p_2) = (x \vee (p_1 \wedge p_2)) - r(p_1 \wedge p_2)$, we have a contradiction as $q \prec x$. Thus, in each of the cases we get a contradiction and consequently we must have $x \wedge y \prec q$. $\square$

Also, since $q_1 \prec z_1$, $q_2 \prec z_2$ and $q \prec x$, by semimodularity we have $q \vee q_2 \prec x \vee z_2$ and $q \vee q_1 \prec x \vee z_1$.

Claim 2.16. $x \wedge y \prec q_1$.

Proof. Suppose there exists p_1 such that $x \wedge y \prec p_1 < q_1$. By semimodularity, we have $q \prec q \vee p_1$, $x \prec x \vee p_1$, $z_2 \prec p_1 \vee z_2$ and $q \vee q_2 \prec q \vee q_2 \vee p_1$. Let $z'_2 = p_1 \vee z_2$ and we have $x \vee z_2 \prec z'_2 \vee x$. Let $x_1 = x \vee p_1$ and we have $lt[x_1 \wedge y, x_1 \vee y] < lt[x \wedge y, x \vee y]$.

Since $z'_2 > z_2$ and $P_{x,y}$ is a dual order ideal, we have $z'_2 \in P_{x,y}$. Thus $r(x \vee z'_2) - r(z'_2) = 1$. Also we have $x_1 \vee y = x \vee y$, $x_1 \vee z_1 = x \vee z_1$ and $x_1 \vee z'_2 = x \vee z'_2$. It follows that $y, z_1, z'_2 \in P_{x_1,y}$ and so $z_1 \wedge z'_2 \in P_{x_1,y}$. Thus $r(x_1 \vee (z_1 \wedge z'_2)) - r(z_1 \wedge z'_2) = 1$. Since $q_1 > z_1 \wedge z'_2 \geq p_1$ and $x_1 \vee (z_1 \wedge z'_2) = x \vee z_1$, we have $r(x_1 \vee (z_1 \wedge z'_2)) - r(z_1 \wedge z'_2) > 1$, a contradiction, and so we must have $x \wedge y \prec q_1$. $\square$




Figure 3.

Claim 2.17. $x \wedge y \prec q_2$.

Proof. Is similar to that of Claim 2.16. $\square$

Claim 2.18. $q_1 \vee q_2 < y$.

Proof. Suppose that $q_1 \vee q_2 = y$, since $x \wedge y \prec q_1, q_2$ implies $x \wedge y = q_1 \wedge q_2$ and by semimodularity, $q_1 \prec y$ and $q_2 \prec y$, which is not true and so we must have $q_1 \vee q_2 < y$. $\square$

Now, $x \wedge y \prec q_1, q_2$ implies $x \wedge y = q_1 \wedge q_2$ and by semimodularity, $q_1 \prec q_1 \vee q_2$ and $q_2 \prec q_1 \vee q_2$. Also, $x \wedge y \prec q, q_1$ implies $x \wedge y = q_1 \wedge q$ and so by semimodularity, $q \prec q \vee q_1$ and $q_1 \prec q \vee q_1$. Similarly, $x \wedge y \prec q, q_2$ implies $x \wedge y = q_2 \wedge q$, so by semimodularity, $q \prec q \vee q_2$ and $q_2 \prec q \vee q_2$. Now, $q_1 \prec q_1 \vee q_2$ and $q_2 \prec q_1 \vee q_2$ implies $q_1 = (q_1 \vee q_2) \wedge z_1$ and $q_2 = (q_1 \vee q_2) \wedge z_2$ and by semimodularity, $z_1 \prec (q_1 \vee q_2) \vee z_1 = y$, $z_2 \prec (q_1 \vee q_2) \vee z_2 = y$ and $q_1 \vee q_2 \prec (q_1 \vee q_2) \vee z_2 = y$.

Claim 2.19. $(q_1 \vee q_2) \vee q < x \vee y$.

Proof. Suppose that $(q_1 \vee q_2) \vee q = x \vee y$. Since $q \wedge (q_1 \vee q_2) \prec q$, by semimodularity we have $q_1 \vee q_2 \prec q \vee (q_1 \vee q_2)$, which is not true as $q_1 \vee q_2 \prec y \prec x \vee y$. Therefore $(q_1 \vee q_2) \vee q < x \vee y$. $\square$

Since $q_1 \prec q \vee q_1$, by semimodularity, $(q_1 \vee q_2) \prec (q_1 \vee q_2) \vee q$. Also, $q_1 \prec q_1 \vee q_2$ implies $q_1 \vee q \prec q_1 \vee q_2 \vee q$, $q_2 \prec q_1 \vee q_2$, which further implies $q_2 \vee q \prec q_1 \vee q_2 \vee q$ and also $q_1 \vee q_2 \prec y$ implies $q_1 \vee q_2 \vee q \prec x \vee z$. Hence, L contains the following cover preserving sublattice.




Figure 4.

□

The following result is due to Teo [8].

Corollary 2.20 ([8]). *A lattice L of finite length is not semimodular if and only if L contains a subpentagon $(a \wedge c, a, b, c, a \vee b)$ with the properties*

- (i) $a \wedge c \prec a, b \prec c \prec a \vee b$, or
- (ii) $a \wedge c \prec a, a \wedge c \prec b, c \prec a \vee b$.




Figure 5.

Corollary 2.21. *Let L be a lattice of finite length. Then L is a pseudomodular lattice if and only if it does not contain a sublattice isomorphic to a cover preserving lattice as depicted in Figure 4 or Figure 5 (a) or Figure 5 (b).*

References

- [1] *G. Birkhoff*: Lattice Theory. Colloquium Publications 25. AMS, Providence, 1967. [zbl](#) [MR](#) [doi](#)
- [2] *A. Björner, L. Lovász*: Pseudomodular lattices and continuous matroids. *Acta Sci. Math.* 51 (1987), 295–308. [zbl](#) [MR](#)
- [3] *A. Dress, W. Hochstättler, W. Kern*: Modular substructures in pseudomodular lattices. *Math. Scand.* 74 (1994), 9–16. [zbl](#) [MR](#) [doi](#)
- [4] *A. Dress, L. Lovász*: On some combinatorial properties of algebraic matroids. *Combinatorica* 7 (1987), 39–48. [zbl](#) [MR](#) [doi](#)
- [5] *G. Grätzer*: General Lattice Theory. Pure and Applied Mathematics 75. Academic Press, New York, 1978. [zbl](#) [MR](#) [doi](#)
- [6] *L. Haskins, S. Gudder*: Heights on posets and graphs. *Discrete Math.* 2 (1972), 357–382. [zbl](#) [MR](#) [doi](#)
- [7] *M. Stern*: Semimodular Lattices: Theory and Applications. Encyclopedia of Mathematics and Its Applications 73. Cambridge University Press, Cambridge, 1999. [zbl](#) [MR](#) [doi](#)
- [8] *K. L. Teo*: Diagrammatic characterizations of semimodular lattices of finite length. *Southeast Asian Bull. Math.* 12 (1988), 135–140. [zbl](#) [MR](#)

Authors' addresses: *Manoj Dhake*, Department of Mathematics, S. P. Pune University, Ganeshkhind Rd, Pune, Maharashtra 411007, India, e-mail: manojdhake2012@gmail.com; *Sachin Ballal* (corresponding author), School of Mathematics and Statistics, University of Hyderabad, Hyderabad Central University Rd, Hyderabad, Telangana, 500046, India, e-mail: sachinballal@uohyd.ac.in; *Vilas Kharat, Rupesh S. Shewale*, Department of Mathematics, S. P. Pune University, Ganeshkhind Rd, Pune, Maharashtra 411007, India, e-mail: laddoo1@yahoo.com, rsshewale@gmail.com.