

Jan Menu

A partition of $\mathbb{R}$ in two homogeneous and homeomorphic parts

In: Zdeněk Frolík (ed.): Abstracta. 5th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1977. pp. 43–49.

Persistent URL: <http://dml.cz/dmlcz/701088>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

A PARTITION OF $\mathbb{R}$ IN TWO HOMOGENEOUS AND HOMEOMORPHIC PARTS

Jan MENU

1. INTRODUCTION.

In this paper we give a sketch of the construction of the partition.

Consider the maps $P(x) = 3^n x + 2k \cdot 3^m$; $n, k, m \in \mathbb{Z}$, and denote by $P = \{P_n | n \in \mathbb{N}\}$ the set of these. In paragraph 2, it is proved that if $A \subset \mathbb{R}$ is stable for P , and $1 \in A$, then A is necessarily homogeneous. We then consider the sets $X(A)$, the orbits of a by P . In paragraph 4 it is proved that if a function f with certain properties exists, the required partition can be constructed. In paragraph 5 a sketch is given of the construction of such a function, omitting the technical parts.

I am indebted to Prof. Maurice to have suggested the problem and for valuable discussions.

2. PROPOSITION. Let $A \subset \mathbb{R}$ be stable for the maps P_n , $1 \in A$, then A is homogeneous.

PROOF. Let $x \in A$, it is sufficient to construct a homeomorphism $h : A \rightarrow A$, such that $h(x) = 1$

a) apply a translation $h_0 : A \rightarrow A$, $x \rightarrow x + 2k \cdot 3^m$ such that $h(x_0) \in]0, 2[$

b) 1) if $h_0(x) \in]\frac{2}{3}, \frac{4}{3}[$, define $h_1 = h_0$

2) if $h_0(x) < \frac{2}{3}$, then there exists just one $n_1 \in \mathbb{Z}$ such that

$$2 \cdot 3^{n_1} < h_0(x) < 2 \cdot 3^{n_1-1}$$

$$\text{define } h_1(y) = \begin{cases} h_0(y) & \text{if } h_0(y) \notin]0, 2[\\ 3^{n_1+1} \cdot h_0(y) & \text{if } 0 < h_0(y) < 2 \cdot 3^{n_1} \\ h_0(y) + (2 - 2 \cdot 3^{n_1}) & \text{if } 2 \cdot 3^{n_1} < h_0(y) < \frac{4}{3} \\ 3^{-n_1-1} (h_0(y) - \frac{4}{3}) + (2 - 2 \cdot 3^{n_1}) & \text{if } \frac{4}{3} < h_0(y) < 2 \end{cases}$$

$$\Rightarrow h_1(x) \in]\frac{2}{3}, \frac{4}{3}[$$

3) if $h_0(x) > \frac{4}{3}$, construct h_1 in an analogous way.

c) Suppose h_n is constructed, $h_n(x) \in]1-3^{-n}, 1+3^{-n}[$

1) if $h_n(x) \in]1-3^{-n-1}, 1+3^{-n-1}[$

$$\text{define } h_{n+1} = h_n$$

2) if $h_n(x) \in]1-3^{-n}, 1-3^{-n-1}[$

then there exists just one $n_{n+1} \in \mathbb{Z}$ such that

$$1-3^{-n} + 2 \cdot 3^{-n+n_{n+1}} < h_n(x) < 1-3^{-n} + 2 \cdot 3^{n_{n+1}-n+1}$$

$$h_{n+1}(y) = \begin{cases} h_n(y) & \text{if } h_n(y) \notin]1-3^{-n}, 1+3^{-n}[\\ 1-3^{-n} + 3^{-n_{n+1}-1} (h_n(y) - (1-3^{-n})) & \text{if } 1-3^{-n} < h_n(y) < 1-3^{-n} + 2 \cdot 3^{n_{n+1}-n} \\ h_n(y) + 2 \cdot 3^{-n-1} - 2 \cdot 3^{n_{n+1}-n} & \text{if } 1-3^{-n} + 2 \cdot 3^{n_{n+1}-n} < h_n(y) < 1+3^{-n-1} \\ 3^{n_{n+1}+1} (h_n(y) - (1+3^{-n-1})) + 1+3^{-n} - 2 \cdot 3^{n_{n+1}-n} & \text{if } 1+3^{-n-1} < h_n(y) < 1+3^{-n} \end{cases}$$

$$\Rightarrow h_{n+1}(y) \in]1-3^{-n-1}, 1+3^{-n-1}[$$

3) if $h_n(x) \in]1+3^{-n-1}, 1+3^{-n}[$, construct h_{n+1} in an analogous way.

d) define $h(y) = \lim_n h_n(y)$.

It is easy to see that h is a homeomorphism of A and $h(x) = 1$.

3. The classes $X(a)$, $a \in \mathbb{R}$.

1. Define $X(a) = \{3^m a + 2k \cdot 3^n \mid m, n, k \in \mathbb{Z}\}$. Every $X(a)$ is countable, and the set $\{X(a) \mid a \in \mathbb{R}\}$ is a partition of $\mathbb{R}$.

$$2. a + 1 \in X(a) \Leftrightarrow a + 1 = 3^n a + 2k \cdot 3^m \quad (n, k, m \in \mathbb{Z})$$

$$\Leftrightarrow a(3^n - 1) = 1 - 2k \cdot 3^m$$

$$\Leftrightarrow a = \frac{1 - 2k \cdot 3^m}{3^n - 1}$$

$$\Leftrightarrow a \in X\left(\frac{2l+1}{3^{p-1}}\right) \text{ for some } p, l \in \mathbb{N} \text{ and } 0 < 2l+1 < 2 \cdot 3^p.$$

In this case $X(a+1) \cap X(a) \neq \emptyset$, and $X(a+1) = X(a)$. Denote by

$(K_n)_n$ these classes.

3. Let $a \in X\left(\frac{2l+1}{3^{n-1}}\right) = K_q$ where $l, n \in \mathbb{N}$, $0 < 2l+1 < 2 \cdot 3^n$ and such that $\forall k', n' \in \mathbb{N}; n' < n$:

$$\frac{2k'+1}{3^{n'}-1} \notin X\left(\frac{2l+1}{3^{n-1}}\right).$$

then there are unique k, p, m with $0 \leq m < n$, such that

$$a = 3^m \cdot \frac{2l+1}{3^{n-1}} + \frac{k}{3^p}.$$

Define $D(q, k) = \left\{ x \mid x = 3^m \cdot \frac{2l+1}{3^{n-1}} + \frac{r}{3^p}, \begin{matrix} 0 \leq p \leq k \\ 0 \leq m < n \\ r \in \mathbb{Z} \end{matrix} \right\}$ where $K_q = X\left(\frac{2l+1}{3^{n-1}}\right)$.

4. $R_1 = \mathbb{R} \setminus X(0)$.

4. THEOREM. Suppose there is a continuous function $f : R_1 \rightarrow R_1$, such that :

$$\left\{ \begin{array}{l} 1) f^2 = 1_{R_1} \\ 2) \forall x : f(x) \notin X(x+1) \\ 3) \forall x : X(f(x)) = f(X(x)) \\ 4) \text{ if } g(x) = f(x) + 1 \text{ then :} \\ \quad \forall x \notin X(1), \forall n \in \mathbb{N} : x \notin X(g^{2n+1}(x)) \\ 5) f(1) \in X(1) \end{array} \right.$$

then there is a partition of R in two homeomorphic parts.

Remark. $g(X(x)) = X(g(x))$, $\forall x \in R_1$.

PROOF of the theorem.

1. $\forall x \in R_1 \setminus X(1)$; define

$$A_x = \cup \{X(g^{2n}(x)) \mid n \in \mathbb{Z}\}$$

$$B_x = \cup \{X(g^{2n+1}(x)) \mid n \in \mathbb{Z}\}.$$

Suppose $g^{2n}(x) \in X(g^{2m+1}(x))$, then $x \in g^{-2n}(X(g^{2m+1}(x))) = X(g^{2m-2n+1}(x))$ which contradicts (4), thus $A_x \cap B_x = \emptyset$.

2. Choose $x_1 \notin X(1) \cup X(0)$. Define $A_1 = A_{x_1}$, $B_1 = B_{x_1}$
 $A_1 \cap B_1 = \emptyset$.

3. Suppose A_β and B_β are defined for every ordinal $\beta < \alpha$, $A_\beta \cap B_\beta = \emptyset$.
 Then a) if α is a limit-ordinal, define

$$\left\{ \begin{array}{l} A_\alpha = \cup \{A_\beta \mid \beta < \alpha\} \\ B_\alpha = \cup \{B_\beta \mid \beta < \alpha\} \end{array} \right.$$

b) if α is not a limit-ordinal,

i) $A_\beta \cup B_\beta \cup X(0) \cup X(1) = \mathbb{R}$, for some $\beta < \alpha$ define

$$A_\alpha = \cup \{A_\beta \mid \beta < \alpha\}$$

$$B_\alpha = \cup \{B_\beta \mid \beta < \alpha\}$$

ii) $A_{\alpha-1} \cup B_{\alpha-1} \cup X(0) \cup X(1) = C_{\alpha-1} \neq \mathbb{R}$. Choose $x_\alpha \in \mathbb{R} \setminus C_{\alpha-1}$.

$$\text{Define } \begin{cases} A_\alpha = A_{\alpha-1} \cup A_{x_\alpha} \\ B_\alpha = B_{\alpha-1} \cup B_{x_\alpha} \end{cases}$$

for every α such that $A_\alpha \cap B_\alpha = \emptyset$ and $g(A_\alpha) = B_\alpha$. There exists

an ordinal α_0 such that $A_{\alpha_0} \cup B_{\alpha_0} \cup X(1) \cup X(0) = \mathbb{R}$.

$$4. \text{ Define } \begin{cases} A = \cup \{A_\alpha \mid \alpha < \alpha_0\} \cup X(1) \\ B = \cup \{B_\alpha \mid \alpha < \alpha_0\} \cup X(0). \end{cases}$$

$$\text{Then : } \begin{cases} A \cap B = \emptyset \\ A \cup B = \mathbb{R} \end{cases}$$

$g : A \rightarrow B$ is a homeomorphism.

4.2. This proves that $\mathbb{R} = A \cup B$, where

$$\begin{cases} A \cap B = \emptyset \\ A \cong B \\ A \text{ and } B \text{ homogeneous.} \end{cases}$$

5. Sketch of the construction of f .

5.1. a) We only consider intervals $[a, b]$ with $a, b \in X(0)$, and

$$b - a = 2 \cdot 3^k \text{ for some } k \in \mathbb{Z}.$$

A net N on such an interval is a finite set of points

$a = a_0 < a_1 < \dots < a_{n-1} < a_n = b$, such that $\forall i < n-1$, there exists a $m \in \mathbb{Z} : a_{i+1} - a_i = 2 \cdot 3^m$:

b) Let N be a net on $[0, 2]$, $x \in [0, 2]$. Denote :

$$V_N(x) = \cup \{V \mid V \in N, x \in \bar{V}\}.$$

c) Let $[a, b]$ be an interval, N a net on $[a, b]$. A function

- $$\left\{ \begin{array}{l} 1) \forall i \leq n-1 : h|_{[a_i, a_{i+1}[} \text{ is a translation onto} \\ \quad \text{some } [a_j, a_{j+1}[, j \leq n-1 \\ 2) h \text{ is the identity on } [a_0, a_1[, [a_{n-1}, a_n[\text{ and on} \\ \quad \text{the interval containing } \frac{a+b}{2} \\ 3) h^2 = 1|_{[a, b] \cap R_1} \\ 4) \forall x \in R_1 : h(x) + 2 = h(x+2) \end{array} \right.$$

5.2. At the n^{th} step is constructed the following :

- 1) A finite set $B_n \subset \cup \{K_m | m \in \mathbb{N}\}$, such that $B_{n-1} \subset B_n$ and $\cup \{B_m | m \in \mathbb{N}\} = \cup \{K_m | m \in \mathbb{N}\}$.
- 2) $\forall k \leq n$ and $x, y \in B_n \cap K_k$ a $P_{x,y}^k \in \{P_m | m \in \mathbb{N}\}$, such that $P_{x,y}^k(x) = y$, and if $P_n \{P_{x,y}^k | x, y \in B_n \cap K_k, k \leq n\}$, then $P_{n-1} \subset P_n$ and P_n is a transitive representation group.
- 3) $\forall x \in B_{n-1} \cap K_k, k \leq n-1$, a $V_{x,n-1}^k \in V(x)$, such that $P_{x,y}^k(V_{x,n-1}^k) = V_{y,n-1}^k$ and if $k' \leq n, 1 \leq n-1, V_{x,n-1}^k \cap V_{y,1}^{k'} \neq \emptyset \Rightarrow V_{x,n-1}^k \subset V_{y,1}^{k'}$. Denote $V_n = \cup \{V_{x,n-1}^k | x \in B_{n-1} \cap K_k, k \leq n-1\}$.
- 4) A net N_{n-1} that refines N_{n-2} and contains the endpoints of the $V_{x,n-1}^k, k \leq n-1, x \in B_{n-1} \cap K_k$, and such that $P_{x,y}^k(N_{n-1}|_{V_{x,n-1}^k}) = N_{n-1}|_{V_{y,n-1}^k}$.
- 5) A function f_{n-1} such that
 - a) $f_{n-1}|_{[0,2] \cap (R_1 \setminus F_{n-2}(V_{n-1}))} = 1$, where $F_{n-2} = f_{n-2} \circ f_{n-3} \circ \dots \circ f_1$.
 - b) f_{n-2} is a N_2 -elementary function on every $F_{n-2}(V_{x,n-1}^k)$

$$c) \forall x, y : P_{x,y}^k \circ f_{n-1} \circ F_{n-2}(x') = f_{n-1} \circ F_{n-2} \circ P_{x,y}^k(x'), \\ x' \in V_{x,n-1}^k.$$

6) Moreover, there is taken care of the following :

$$a) x \in K_{2m-1} \Rightarrow \lim_n F_n(x) \in K_{2m}$$

b) the condition 4) is satisfied in the n^{th} step outside of a set of length $< 3^{-n}$.

Define $f = \lim_n F_n$, then clearly f has the required properties, and this completes the proof.