

Edward Grzegorek

On Ulam's problem on families of measures

In: Zdeněk Frolík (ed.): Abstracta. 6th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1978. pp. 45–48.

Persistent URL: <http://dml.cz/dmlcz/701119>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1978

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

SIXTH WINTER SCHOOL (1978)

ON ULAM'S PROBLEM ON FAMILIES OF MEASURES

by

E. GRZEGOREK

Throughout, $|S|$ denotes the cardinality of the set S , $\mathcal{P}(S)$ the power set of S , $[S]^\gamma = \{X \subset S : |X| = \gamma\}$, and $V=L$ denotes Gödel's axiom of constructability. Small greek letters denote ordinals, with $\aleph, \mu$ always denoting infinite cardinals and λ, ν any (finite or infinite) cardinals. The following corollary follows from our Theorem 3.

COROLLARY. Let F be a family of σ -fields of subsets of the real line S , such that $[S]^\lambda \subset A$ and $A \neq \mathcal{P}(S)$ for every $A \in F$. Then

- a) $|F| < \omega$ implies $\bigcup F \neq \mathcal{P}(S)$;
- b) If $2^\omega = \omega_1$ then $|F| \leq \omega$ implies $\bigcup F \neq \mathcal{P}(S)$;
- c) If $V = L$ then $|F| \leq \omega_1$ implies $\bigcup F \neq \mathcal{P}(S)$.

The Corollary can be strengthened even under weaker set theoretical assumption (see Theorem 3). In the case of an additional assumption that on each $A \in F$ it is possible to define a non-trivial measure (or even that A satisfies only certain chain condition), the Corollary has been known. In that case, a) is due to Ulam (see [1]), b) is a theorem of Alaoglu - Erdős (see [1] and also [4] and [3]), and c) is a theorem of Prikry (see [4], for generalizations see [3], for strengthenings and further generalizations see [6]. In case on each $A \in F$ it is possible to define a non-trivial two-valued measure, c) is a theorem of Jensen (see [0]).

The strongest and the most general results connected with a problem of Ulam on families of measures (see problem 81 of [2] and also [8]) have been recently obtained by Taylor in [6]. The main subject of this note is a generalization of two theorems of Taylor in [6].

If $Q \subset \mathcal{P}(S)$ then we define $I(Q) = \{X \in Q : \mathcal{P}(X) \subset Q\}$. Q will be called μ -complete iff for every $X \subset Q$ such that $|X| < \mu$ we have $\bigcup X \in Q$. Remark that if Q is μ -complete then $I(Q)$ is a μ -complete ideal on S . Q will be called non-trivial iff $[S]^\lambda \subset Q$ and $Q \neq \mathcal{P}(S)$.

A family $F \subset \mathcal{P}(\mathcal{P}(S))$ will be called γ -saturated

w.r.t. I , where I is an ideal on S such that $I \subset \bigcap \{I(A) : A \in F\}$, iff for every collection $\{X_\alpha : \alpha < \gamma\} \subset \mathcal{P}(S) - \mathcal{U}_F$ there exists $\{\alpha, \beta\} \in [\gamma]^2$ such that $X_\alpha \cap X_\beta \notin I$.

A family $F \subset \mathcal{P}(\mathcal{P}(S))$ will be called γ -saturated iff F is γ -saturated w.r.t. $I = \bigcap \{I(A) : A \in F\}$.

The following two definitions are central for the considerations of this note.

If $Q \subset \mathcal{P}(\mathcal{P}(\kappa))$ then the symbol

$$\langle \kappa : \lambda, \mu \rangle \xrightarrow{Q} \gamma$$

denotes the following assertion.

If $F \subset Q$, $|F| \leq \lambda$ and $I(A)$ is μ -complete for every $A \in F$ then F is not γ -saturated.

If $Q \subset \mathcal{P}(\mathcal{P}(\kappa))$ and I is an ideal on κ (we do not exclude the case $I = \{\emptyset\}$) then the symbol

$$\langle \kappa : \lambda, \mu \rangle \xrightarrow{Q} \langle \gamma, I \rangle$$

denotes the following assertion.

If $F \subset Q$, $|F| \leq \lambda$, $I \subset \bigcap \{I(A) : A \in F\}$ and $I(B)$ is μ -complete for every $B \in F$ then F is not γ -saturated w.r.t. I .

In case Q is a set of all non-trivial ideals on κ the notation $\langle \kappa : \lambda, \mu \rangle \xrightarrow{Q} \gamma$ was introduced by Taylor in [6]. If Q is a set of all non-trivial ideals on κ then instead of $\langle \kappa : \lambda, \mu \rangle \xrightarrow{Q} \gamma$ and $\langle \kappa : \lambda, \mu \rangle \xrightarrow{Q} \langle \gamma, I \rangle$ we will write $\langle \kappa : \lambda, \mu \rangle \rightarrow \gamma$ and $\langle \kappa : \lambda, \mu \rangle \rightarrow \langle \gamma, I \rangle$, respectively (i.e. we suppress the superscript Q in this case).

For a fixed cardinal κ we define

$$R = \{A \subset \mathcal{P}(\kappa) : A \text{ is non-trivial and } \forall (a \in A) \forall (b \in A) (a \cap b \in A \text{ and } a - b \in A)\}$$

We have the following theorem.

THEOREM 1. Assume $\lambda \leq \gamma \leq \omega$. Then we have

a) If I is a $(\lambda + \omega)$ -complete ideal on κ then

$$\langle \kappa : \lambda, \mu \rangle \xrightarrow{R} \langle \gamma, I \rangle \text{ iff } \langle \kappa : \lambda, \mu \rangle \rightarrow \langle \gamma, I \rangle.$$

a') If $\lambda \leq \mu$ then

$$\langle \kappa : \lambda, \mu \rangle \xrightarrow{R} \gamma \text{ iff } \langle \kappa : \lambda, \mu \rangle \rightarrow \gamma.$$

From Theorem 1 we have in particular the following result:

$$\langle \omega_1 : \omega_1, \omega_1 \rangle \rightarrow \omega_1 \text{ iff } \langle \omega_1 : \omega_1, \omega_1 \rangle \xrightarrow{R} \omega_1.$$

This (and also our Theorem 3) should be compared with the comments of the authors of [2] on the problem 81 of Ulam

(see also [8]).

With the help of Theorem 1 we will generalize the following results of Taylor (Theorem 2.2 and Theorem 4.4. of [6]). We formulate them in a little more general form, which easily follows from the original one.

THEOREM 2 (TAYLOR)

a) Assume $\nu \geq \lambda^+ + \omega$, $\mu \geq \lambda^+ + \omega$, $\lambda < \kappa$ and I is a $(\lambda^+ + \omega)$ -complete ideal on κ . Then

$$\langle \kappa: \lambda, \mu \rangle \rightarrow \langle \nu, I \rangle \quad \text{iff} \quad \langle \kappa: 1, \mu \rangle \rightarrow \langle \nu, I \rangle$$

$$b) \langle \omega_1: \omega_1, \omega_1 \rangle \rightarrow \omega_2 \quad \text{iff} \quad \langle \omega_1: 1, \omega_1 \rangle \rightarrow \langle \omega_2, [\omega_1]^{<\omega_1} \rangle$$

Recall that the above theorem of Taylor is a strengthening and a generalization of results of Ulam, Alaoglu - Erdős (see [1]), Jensen (see [0]), Prikry (see [4]) and of the present author (see [3]). By Theorem 1 and Theorem 2 we have the following generalization of Theorem 2.

THEOREM 3. a) Assume I is a $(\lambda^+ + \omega)$ -complete ideal on κ and $\nu \geq (\lambda^+ + \omega)$, $\mu \geq \lambda^+ + \omega$, $\lambda < \kappa$. Then

$$\langle \kappa: \lambda, \mu \rangle \xrightarrow{R} \langle \nu, I \rangle \quad \text{iff} \quad \langle \kappa: 1, \mu \rangle \rightarrow \langle \nu, I \rangle.$$

$$b) \langle \omega_1: \omega_1, \omega_1 \rangle \xrightarrow{R} \omega_2 \quad \text{iff} \quad \langle \omega_1: 1, \omega_1 \rangle \rightarrow \langle \omega_2, [\omega_1]^{<\omega_1} \rangle.$$

Remark that if we replace R by $R_0 \subset R$, where R_0 is a collection of families of subset of κ , satisfying certain natural chain conditions, then Theorem 3 becomes a known result which easily follows directly from Theorem 2 (see Corollary 4.13 of [6], compare also [3] and [4]).

To see for which $\kappa, \lambda, \mu, \nu$ Theorem 3 works, recall the following well known facts. $\langle \kappa^+: 1, \kappa^+ \rangle \rightarrow \kappa^+$ and $\langle 2^\kappa: 1, \kappa^+ \rangle \rightarrow \omega$ holds for every κ (see [7]). $\langle \kappa: 1, \mu \rangle \rightarrow \mu$ holds for every κ which is less than the first weakly inaccessible cardinal and every $\mu \leq \kappa$ (easily follows from the first previous relations). $\langle \kappa: 1, \omega_1 \rangle \rightarrow \omega$ holds for every κ which is less than the first strongly inaccessible cardinal (see [7]). By results of Tarski and Solovay the relations holds if κ is even larger. It is also well known that the axiom of constructability ($V = L$) implies $\langle \omega_1: 1, \omega_1 \rangle \rightarrow \langle \omega_2, [\omega_1]^{<\omega_1} \rangle$ (see [5]).

The elementary proof of Theorem 1 will be submitted elsewhere.

References:

- [0] K. Devlin, Aspects of constructibility, Lecture notes in math. Vol. 354.
- [1] P. Erdős, Some remarks on set theory, Proc. Amer. Math. Soc. 1(1950), 127-141.
- [2] P. Erdős, A. Hajnal, Unsolved problems in set theory, Proc. Symp. Pure Math. 13, Part I. 1971, ed. D. Scott, 17-48.
- [3] E. Grzegorek, A remark on a paper by Karel Prikry "Kurepa's hypothesis and a problem of Ulam on families measures", (submitted to Colloq. Math.).
- [4] K. Prikry, Kurepa's hypothesis and a problem of Ulam on families of measures, Monatshefte für Math. 81(1976), 41-57.
- [5] -, Kurepa's hypothesis and $\aleph_1$ -complete ideals, Proc. Amer. Math. Soc. 38(1973), 617-620.
- [6] A.D. Taylor, On saturated sets of ideals and Ulam's problem, (preprint).
- [7] S.M. Ulam, Zur Masstheorie in der allgemeinen Mengenlehre, Fund. Math. 16(1930), 140-150.
- [8] -, A collection of mathematical problems, Interscience Publishers, Inc., New York, 1960.