

Gliceria Godini

On minimal points with respect to a set in Banach spaces

In: Zdeněk Frolík (ed.): Abstracta. 8th Winter School on Abstract Analysis.
Czechoslovak Academy of Sciences, Praha, 1980. pp. 82–85.

Persistent URL: <http://dml.cz/dmlcz/701182>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic,
1980

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project
DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

8th WINTER SCHOOL ON ABSTRACT ANALYSIS (1980)

ON MINIMAL POINTS WITH RESPECT TO A SET IN BANACH SPACES

by

G. GODINI

Let Y be a linear subspace of the (real) normed linear space X and $\lambda \geq 1$. We assign to each set $M \subset Y$ a set $M_{Y,X}^\lambda \subset X$ in the following way: $x \in M_{Y,X}^\lambda$ if there exists no $y \in Y$, $y \neq x$, such that:

$$\|y - m\| \leq \lambda \|x - m\| \quad \text{for each } m \in M.$$

When $Y = X$ and $\lambda = 1$ then $M_{X,X}^1$ is the set of minimal points with respect to M studied by B. Beauzamy and B. Maurey in [1] and denoted there by $\min M$. When $Y \subset X$ and $\lambda = 1$ then $M_{Y,X}^1$ was introduced and studied in [2].

For each $M \subset Y$ and $1 \leq \lambda \leq \mu$ we have $M_{Y,Y}^\lambda = M_{Y,X}^\lambda \cap Y$, $M_{X,X}^\lambda \subset M_{Y,X}^\lambda$ and $M_{Y,X}^\mu \subset M_{Y,X}^\lambda$, the inclusions above (as well as $M_{Y,Y}^\lambda \subset M_{Y,X}^\lambda$) being strictly in general, as examples show.

B. Beauzamy and B. Maurey [1] proved the following result: Let X be a reflexive, strictly convex and smooth Banach space and Y a closed linear subspace of X . If $\min Y = Y$ (in our notation $Y_{X,X}^1 = Y$) then there exists a (unique) norm one linear projection of X onto Y . They also remarked that the existence of a norm one linear projection of X onto Y implies $\min Y = Y$. In [2] we gave also a necessary and sufficient condition for the existence of a norm one linear projection of X onto Y , weakening the conditions on X (requiring only the smoothness of X) but strengthening the condition $\min Y = Y$.

When there are no restrictions on X we have the following result. Let $\tilde{\mathcal{P}}(X, Y, \lambda)$ be the set of all mappings $\tilde{P}: X \rightarrow Y$ with the following properties:

- (1) $\tilde{P}^2(x) = \tilde{P}(x)$ ($x \in X$)
- (2) $\tilde{P}(\alpha x) = \alpha \tilde{P}(x)$ ($x \in X, \alpha \in \mathbb{R}$)
- (3) $\tilde{P}(x+y) = \tilde{P}(x) + \tilde{P}(y)$ ($x \in X, y \in Y$)
- (4) $\|\tilde{P}(x)\| \leq \lambda \|x\|$ ($x \in X$)

Theorem 1. Let Y be a closed linear subspace of the normed linear space X and $\lambda \geq 1$. We have $\tilde{\mathcal{P}}(X, Y, \lambda) \neq \emptyset$ if and only if $Y_{Y, X}^\lambda = Y$.

Let us denote $S_X = \{x \in X : \|x\| = 1\}$, $\text{sm } S_X$ the set of all $x \in S_X$ such that there exists a unique $x_X^* \in S_X$ with $x_X^*(x) = 1$, and $\mathcal{P}(X, Y, \lambda)$ the set of all linear projections P of X onto Y with $\|P\| \leq \lambda$.

Theorem 2. ([2]). Let X be a normed linear space and Y a closed linear subspace of X . A necessary, and if $S_Y \subset \text{sm } S_X$ also sufficient condition for $\mathcal{P}(X, Y, 1) \neq \emptyset$ is that $Y_{Y, X}^1 = Y$. If $S_Y \subset \text{sm } S_X$ then $\mathcal{P}(X, Y, 1)$ contains at most one element.

When $\lambda \geq 2$ we can exhibit subspaces $Y \subset X$ with $\tilde{\mathcal{P}}(X, Y, \lambda) \neq \emptyset$. Indeed, for $Y \subset X$ and $x \in X$ let $P_Y(x)$ be the set of all best approximations of x out of Y , i.e., $P_Y(x) = \{y_0 \in Y : \|x - y_0\| = \text{dist}(x, Y)\}$. Y is called a proximal subspace of X if $P_Y(x) \neq \emptyset$ for each $x \in X$ and a Čebyšev subspace of X if $P_Y(x)$ contains exactly one element for each $x \in X$. We shall denote the elements of $P_Y(x)$ by $\pi_Y(x)$. When Y is a proximal subspace of X we can choose a selection $\pi_Y(x) \in P_Y(x)$ satisfying the conditions (1)-(3), and since $\|\pi_Y(x)\| \leq 2\|x\|$ for

each $x \in X$, it follows that $\tilde{\mathcal{P}}(X, Y, \lambda) \neq \emptyset$ for each $\lambda \geq 2$, and so $Y_{Y, X}^\lambda = Y$ for each $\lambda \geq 2$. Using this remark and a now classical result of J. Lindenstrauss and L. Tzafriri [3] it follows that for $\lambda \geq 2$ we have not a result similar with Theorem 2.

Some immediate consequences of the above results are:

Corollary 1. If $S_Y \subset \text{sm } S_X$ then $\tilde{\mathcal{P}}(X, Y, 1) = \mathcal{P}(X, Y, 1)$ and $\tilde{\mathcal{P}}(X, Y, 1)$ contains at most one element.

Corollary 2. Let Y be a proximal subspace of the normed linear space X such that $S_Y \subset \text{sm } S_X$ and $\|\pi_Y(x)\| \leq \|x\|$ for each $\pi_Y(x) \in P_Y(x)$ and each $x \in X$. Then Y is a Čebyšev subspace of X and π_Y is linear. Moreover $\mathcal{P}(X, Y, 1) = \{\pi_Y\}$.

We conclude this note with the following particular case. Let E be a normed linear space and we regard E as a subspace of its second dual E^{**} . If $X = E^{**}$ and $Y = E$, then for each $M \subset E$ we denote $\text{MIN } M = M_{E, E^{**}}^1$. F. Sullivan [4] called a Banach space E to be very smooth if $S_E \subset \text{sm } S_{E^{**}}$. Examples of very smooth nonreflexive spaces as well as some properties of very smooth spaces are given in [4]. The last part of Theorem 2 is a generalization of the following result of [4]. If E is a very smooth Banach space then $\mathcal{P}(E^{**}, E, 1)$ contains at most one element. An immediate consequence of Theorem 2 is:

Corollary 3. ([2]). If E is a very smooth Banach space, then $\mathcal{P}(E^{**}, E, 1) \neq \emptyset$ if and only if $\text{MIN } E = E$.

The proofs of our results which are not contained in [2] will be given elsewhere in a more general setting.

References

- [1] B. Beauzamy et B. Maurey : Points Minimaux et Ensembles Optimaux dans les Espaces de Banach, J.Functional Analysis, 24(1977), 107-139.
- [2] G. Godini : On minimal points, (to appear).
- [3] J. Lindenstrauss and L. Tzafriri : On complemented subspace problem, Israel J.Math., 9(1971), 263-269.
- [4] F. Sullivan : Geometrical properties determined by the higher duals of a Banach space, Illinois J. of Math., 21(1977), 315-331.