

Cosimo Guido

Connected spaces which are not strongly connected

In: Zdeněk Frolík (ed.): Proceedings of the 12th Winter School on Abstract Analysis, Section of Topology. Circolo Matematico di Palermo, Palermo, 1984. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 6. pp. [149]–153.

Persistent URL: <http://dml.cz/dmlcz/701836>

Terms of use:

© Circolo Matematico di Palermo, 1984

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

CONNECTED SPACES WHICH ARE NOT STRONGLY CONNECTED (*)

Cosimo Guido

Sunto. In questa nota vengono individuate due classi di spazi topologici, rispettivamente T_0 non- T_1 e T_1 non- T_2 , che sono connessi ma non fortemente connessi.

INTRODUCTION. We recall that a topological space (S, τ) is *maximal connected* if it is connected and no connected topology τ' on S exists which is strictly finer than τ , (S, τ) is *strongly connected* if τ is coarser than a maximal connected topology τ' on S .

The existence of maximal connected spaces verifying some separation axioms has been often investigated; until now it is an open question whether a regular maximal connected space exists or not.

Any-way every maximal connected space is a T_0 space.

We remark that each connected topology on a finite set is strongly connected.

Now if we consider only T_0 topological spaces with infinitely many points, the following questions can be asked.

Do there exist T_0 non- T_1 strongly connected spaces which have, respectively, maximal connected T_2 or maximal connected T_1 or only maximal connected T_0 expansions?

Do there exist T_1 non- T_2 strongly connected spaces which have, respectively, maximal connected T_2 or only maximal connected T_1 expansions?

Do there exist connected non-strongly connected spaces which are respectively T_2, T_1 non- T_2, T_0 non- T_1 ?

The answers to the first two questions are affirmative by the well-known existence of T_0 non- T_1, T_1 non- T_2 , and T_2 maximal connected spaces.

In some papers, listed below, sufficiently large classes of spaces were pointed out in order to give examples of such spaces.

(*) Subject classification AMS 1980 54A10 54D05 54D10.

Partially supported by a grant of M.P.I.

Guthrie and Stone [5] and Baggs [1] found T_2 connected spaces which are not strongly connected.

In this note we show how to construct T_1 non- T_2 or T_0 non- T_1 connected topological spaces which are not strongly connected.

If (S, τ) is a topological space, we shall denote by $\tau(x)$ the family of open neighbourhoods of x while $\mathcal{T}_\tau(x)$ will denote the neighbourhood filter of x in τ . $\tau|_X$ will be the induced topology on the

subset $X \subseteq S$. If X, Y are subsets of S , $X \setminus Y$ will denote their difference. If, finally, X is a disconnected subspace of (S, τ) , we shall say that A, B divide X in τ if A, B are non-empty open sets of $\tau|_X$

and $A \cup B = X$, $A \cap B = \emptyset$

See [3] for further notations not mentioned here.

1. Let (N, τ_0) be a T_2 connected space with a dispersion point $x_0 \in N$. Choose a point $x \neq x_0$ in N and a non-principal open ultrafilter ν on N containing the family $\{V \setminus \{x\} / V \in \tau_0(x)\}$.

Consider the topology τ on N defined by

$$x \notin A \Rightarrow A \in \tau_0$$

$$A \subseteq N, A \in \tau \iff$$

$$x \in A \Rightarrow A \setminus \{x\} \in \nu \cap \tau_0$$

Trivially (N, τ) is Hausdorff and x_0 is a dispersion point of (N, τ) too. Furthermore we have the following.

LEMMA 1. $U \subseteq N$, $U \cup \{x\} \Rightarrow (N \setminus U) \cup \{x\} \in \mathcal{T}_\tau(x)$.

Proof. Let U be a subset of N . If $U \in \nu$, then $A \in \nu \cap \tau_0$, $A \subseteq U$ exists. $A \cup \{x\}$ is open in τ and $U \cup \{x\} \in \mathcal{T}_\tau(x)$.

If $U \notin \nu$, then $B \in \nu \cap \tau_0$, $B \subseteq (N \setminus U)$ exists hence $B \cup \{x\} \in \tau$ and $(N \setminus U) \cup \{x\} \in \mathcal{T}_\tau(x)$.

LEMMA 2. (N, τ) is connected.

Proof. If (N, τ) were disconnected and A, B divided S in τ and $x \in A$, then we should have

$$B \in \tau_0 \quad \text{and} \quad A \setminus \{x\} \in \nu \cap \tau_0$$

hence B and $A \setminus \{x\}$ would divide $N \setminus \{x\}$ in τ_0 and consequently x would be a cut point of (N, τ_0) .

So we have the assertion since the dispersion point x_0 is the only cut point of (N, τ_0) .

Now consider a point $y \notin N$ and put $S = N \cup \{y\}$. Let σ be the topology on S defined by

$$A \subseteq S, A \in \sigma \iff \begin{array}{l} y \notin A \Rightarrow A \in \tau \\ y \in A \Rightarrow A \setminus \{y\} \in \tau_0 \cap \nu \end{array}$$

Of course $\sigma|_N = \tau$ and (S, σ) is a T_1 non- T_2 space.

Furthermore (S, σ) is connected; indeed if A, B divided S in σ and $y \in A$, then $A \setminus \{y\}$ and B would divide N in τ_0 .

Eventually it can be proved the following lemma.

LEMMA 3. N remains connected in every connected expansion of (S, σ)

Proof. Let $(S, \tau'), \tau' \supseteq \sigma$, be a connected expansion of (S, σ) .

If N were disconnected and X, Y divided N in τ' , where $X = A \cap N$, $Y = B \cap N$ with $A, B \in \tau'$, we should have $y \in A \cup B$ whence $A, B \setminus \{y\}$ or A, B divide S in τ' which contradicts the assumptions.

Then let us consider the case $y \notin A \cup B$ and consequently assume $X = A \in \tau'$, $Y = B \in \tau'$; by lemma 1 we could suppose $X \cup \{x\} \in \mathcal{T}_\tau(x)$ and then $X \cup \{y\}$ would be open in $\sigma \subseteq \tau'$. $X \cup \{y\}$ and Y would divide S in τ' and, again, (S, τ') would be disconnected.

The following theorem can now be easily proved.

THEOREM 1. (S, σ) is not a strongly connected space.

Proof. Otherwise, the connected subspace N of some maximal connected expansion (S, τ') would be maximal connected and (N, τ_0) would be strongly connected which is absurd (see [5] theorem 15).

2. Let (M, τ_1) be a T_1 maximal connected topological space such that the non-empty open sets form an ultrafilter $\mathcal{U} = \tau_1 \setminus \{\emptyset\}$.

Take two points $x \in M$ and $y \notin M$, put $X = M \cup \{y\}$ and consider the topology τ on X defined by

$$\begin{aligned} A \setminus \{y\} \in \tau_1 \quad \text{and} \\ A \subseteq X, A \in \tau \iff \\ y \in A \implies x \in A. \end{aligned}$$

(X, τ) is a T_0 non- T_1 connected space and it is not maximal connected (in fact its proper expansion $A \in \tau' \iff A \setminus \{y\} \in \mathcal{U}$ or $A = \emptyset$ is connected).

Furthermore the following results hold for such a topology.

LEMMA 4. If $\tau' \supseteq \tau$ is a maximal connected non- T_1 topology on X , then M is connected in τ' .

Proof. Let M be disconnected and $A \cap M, B \cap M$ divide M ; $A, B \in \tau'$; suppose $x \in A$, then $x \notin B$ and $y \notin B$ since $\tau'(y) \subseteq \tau'(x)$.

If now $y \in A$, then A, B divide X in τ' .

If $y \notin A$ and $A \in \tau_1$ we must have $A \cup \{y\} \in \tau \subseteq \tau'$ whence $A \cup \{y\}, B$ divide X in τ' ; in the same way $A, B \cup \{y\}$ divide X in τ' if $y \notin A$ and $B \in \tau_1$.

Anyway (X, τ') must be disconnected, a contradiction.

LEMMA 5. If $\tau' \supset \tau$ is a maximal connected topology on X and M is connected in τ' , then τ' is a T_1 topology.

Proof. It follows from the assumption that $(M, \tau'|_M)$ is maximal connected since it is a connected subspace of a maximal connected space; on the other hand $\tau_1 = \tau|_M \subseteq \tau'|_M$ is a maximal connected topology too, so we have $\tau'|_M = \tau_1$.

Consider now $U \in \tau' \setminus \tau$; trivially $U \cap M = U \setminus \{y\} \in \tau_1$ and $y \in U$: furthermore it follows from $U \setminus \{y\} \in \tau_1$, $y \in U$ and $U \notin \tau$ that $x \notin U$. U is actually a neighbourhood of y , in τ' , which does not contain x and consequently τ' is a T_1 topology.

Let now $\sigma \supseteq \tau$ be a T_0 non- T_1 connected topology on X which is T_1 -disconnected, i.e. the least T_1 topology containing σ is disconnected (see [3]).

We are now able to prove the concluding result.

THEOREM 2. (X, σ) is not strongly connected.

Proof. If $\tau' \supseteq \sigma$ were a maximal connected expansion of σ one would have $\tau' \supseteq \sigma \supseteq \tau$; on the other hand τ' would not be T_1 , hence $\tau'|_M$ would be connected, by lemma 4, which contradicts the assumption that τ' is maximal connected, by lemma 5.

REFERENCES

- [1] BAGGS I. "A connected Hausdorff space which is not contained in a maximal connected space", Pacific J. Math. 51 (1974), 11-18.
- [2] EL'KIN A.G. "Maximal connected Hausdorff spaces", Math. Notes 26 (1979) n.5-6, 974-978, 1980.
- [3] GUIDO C. "On strongly connected T_0 spaces", Proceedings of XI Winter School on Abstract Analysis, Suppl. Rend. Circ. Mat. Palermo, 1983.
- [4] GUTHRIE J.A., REYNOLDS D.F., STONE H.E. "Connected expansions of topologies", Bull. Austral. Math. Soc. 9 (1973), 259-265.
- [5] GUTHRIE J.A., STONE H.E. "Spaces whose connected expansions preserve connected subsets" Fund. Math. 80 (1973), 91-100.

- [6] GUTHRIE J.A., STONE H.E., WAGE M.L., "Maximal connected expansions of the reals", Proc. Amer. Math. Soc. 69 (1978), 159-165.
- [7] SIMON P. "An example of maximal connected Hausdorff space", Fund. Math. 100 (1978), n.2, 157-163.
- [8] THOMAS J.P. "Maximal connected topologies", J.Austral. Math.Soc. 8 (1968), 700-705.
- [9] THOMAS J.P. "Maximal connected Hausdorff spaces", Pacific J.Math. 57 (1975), 581-583.

DIPARTIMENTO DI MATEMATICA
UNIVERSITA'
VIA ARNESANO
73100 LECCE (ITALIA)