# Josef Kolomý

Set-valued mappings and structure of Banach spaces

In: Zdeněk Frolík and Vladimír Souček and Marián J. Fabián (eds.): Proceedings of the 14th Winter School on Abstract Analysis. Circolo Matematico di Palermo, Palermo, 1987. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 14. pp. [345]–351.

Persistent URL: http://dml.cz/dmlcz/701907

## Terms of use:

© Circolo Matematico di Palermo, 1987

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

### Josef Kolomý

The theory of monotone (maximal monotone), accretive (maximal accretive) single-valued and multi-valued mappings, intensively studied in the last period, has fruitful applications in the theory of nonlinear partial and ordinary differential and integral equations ([2],[4],[8],[21]).

The aim of this note is to present some known recent results concerning single-valuedness and continuity properties of maximal monotone and the new ones of maximal accretive multivalued mappings and the structure of Banach spaces.

#### 1. Notions and notations.

Let X be a real Banach space,  $X^{\times}$  its dual,  $\angle$ , > the pairing between X and  $X^{\times}$ ,  $S_1(0)$  the unit sphere of X. We shall say that a Banach space X is: (i) smooth if its norm is Gateaux differentiable on  $S_1(0)$ ; (ii) Fréchet smooth if its norm is Fréchet differentiable on  $S_1(0)$ ; (iii) an Asplund space (a week Asplund space) if each convex continuous functional f on X is Fréchet (Gateaux) differentiable on a dense  $G_{\delta}$  subset of X, (iv) an (H)-space, if X is rotund and the following condition is satisfied: if  $(x_n), x \in X$ ,  $x_n \to x$  weakly in X,  $\|x_n\| \to \|x\|$ , then  $x_n \to x$  in the norm of X.

The notions of rotundity (R), local uniform rotundity (LUR), uniform rotundity (UR) of X are used in the usual sense ([15]). Let X,Y be topological spaces,  $T:X \to 2^Y$  a multivalued mapping,  $D(T) = \{u \in X: T(u) \neq \emptyset\}$  its domain,  $G(T) = \{(u,v) \in X \times Y: u \in D(T), v \in T(u)\}$  its graph in the space X x Y . We shall say that T is : (i) upper semi-continuous at  $u_0 \in D(T)$  if for each open subset W of Y such that  $T(u_0) \in W$ , there exists an open neighborhood U of  $u_0$  such that  $T(U) \in W$ ; (ii) lower semicontinuous at  $u_0 \in D(T)$  if for each open subset W of Y such that  $T(u_0) \cap W \neq \emptyset$  there exists an open neighborhood U of  $u_0$  such that  $T(u_0) \cap W \neq \emptyset$  there exists an open neighborhood U of  $u_0$  such that  $T(u) \cap W \neq \emptyset$  for all  $u \in U$ . Let X be a real norhood U of  $u_0$  such that  $T(u) \cap W \neq \emptyset$  for all  $u \in U$ . Let X be a real norhood

This paper is in final form and no version of it will be submitted for publication elsewhere.

med linear space. A mapping  $T:X \longrightarrow 2^{X^{\frac{1}{N}}}$  is said to be: (i) monotone on D(T) if  $\langle u^{\frac{1}{N}} - v^{\frac{1}{N}}, u - v \rangle \stackrel{?}{=} 0$  for each  $u,v \in D(T)$ ,  $u^{\frac{1}{N}} \in T(u)$ ,  $v^{\frac{1}{N}} \in T(v)$ ; (ii) maximal monotone on D(T) if T is monotone on D(T), and its graph G(T) is not properly contained in the graph of any other monotone map.

Now we give some well-known examples of maximal monotone operators.

 $1^{\circ}$ .Let X be a Banach space, f a continuous convex function on X . Then the subdifferential map

2°. If  $T:X \to X^{\mathbb{R}}$  is linear with D(T) = X and  $\langle T(u), u \rangle \stackrel{>}{=} 0$  for each  $u \in X$ , then T is maximal monotone.

Let X be a reflexive Banach space,  $T: X \supset D(T) \longrightarrow X^X$  a closed linear and monotone mapping such that  $\overline{D(T)} = X$ . Then T is maximal monotone if and only if  $T^X$  is monotone. If X is reflexive,  $T: X \longrightarrow 2^{X^X}$  is monotone with  $D(T) \subset X$ , then T is maximal monotone if and only if  $(T + J) X = X^X$  (£21]). The following result (£21]) is useful in applications:

Let X be a reflexive real Banach space,  $T:X \longrightarrow 2^{X^{\frac{1}{2}}}$  a coercive maximal monotone operator on  $D(T)\subset X$ . Then  $T(X)=X^{\frac{1}{2}}$ . For the further results and examples concerning the maximal monotone operators see [2]. [4] and [21].

2. Single-valuedness and continuity properties of maximal monotone multivalued mappings.

Single-valuedness and continuity properties of monotone operators have been studied by Zarantonello [24], Kenderov [19], [20], Fabián [10], [11], Fitzpatrick [12], [13], Zajíček[23], Christensen and Kenderov [6], [7], Jayne and Rogers [18] and others. We recall here only some results which are related to those stated later concerning the accretive multivalued mappings.

Theorem 1 (1191). Let X be a Banach space which admits an equivalent norm such that its dual norm is (R) in  $X^{*}$ . If T:X  $\longrightarrow$   $2^{X^{*}}$  is maximal monotone with D(T) = X, then T is single-valued on a dense  $G_{\bar{g}}$  subset of X.

If X satisfies the renorming condition of the above theorem, then X is a weak Asplund space ([19]). In particular, each WCG (and hence each separable Banach space) is a weak Asplund space.

Theorem 2 ([20]). X is an Asplund space if and only if each maximal monotone mapping  $T:X \longrightarrow 2^{X^{**}}$  with int  $D(T) \neq \emptyset$  is single-valued and upper semicontinuous (with respect to the norm topologies of X and  $X^{**}$ ) on a dense  $G_{\hat{X}}$  subset of X.

It was proved in [6] that the similar result of Theorem 2 holds even in the case when maximal monotonicity of T is replaced by the condition that T is weak compact valued and upper semicontinuous on int D(T) from the norm topology of X into the weak topology of  $X^{M}$ .

Theorem 3 ([9]). Let X be a Banach space such that  $X^{\mathbb{H}}$  is (R) and (H)-space,  $T:X \longrightarrow 2^{X^{\mathbb{H}}}$  a maximal monotone mapping such that int  $D(T) \neq \emptyset$ . Then: (i) there exists a unique lower selection  $T_0$  of T; (ii) for each  $x \in \text{int } D(T)$  at which  $T_0$  is continuous, T(x) is a singleton and T is upper semicontinuous (with respect to the norm topologies of X and  $X^{\mathbb{H}}$ ) at x; (iii) the set  $C(T_0)$  of all those points at which  $T_0$  is continuous is a dense  $G_{\mathcal{F}}$  subset of int D(T).

According to [22] a subset A<X is said to be an  $\alpha$ -angle porous ( $\alpha > 0$ ) if for each  $x \in A$  and each  $\epsilon > 0$  there exist  $z \in B_{\epsilon}(x) = \{u \in X : ||u - x|| < \epsilon \}$  and  $x^{\frac{M}{\epsilon}} \in X^{\frac{M}{\epsilon}}$  such that

An  $\{y \in X : \langle y - z, x^{*} \rangle > \alpha \|x^{*}\| \cdot \|y - z\| \} = \emptyset$ . We shall say that A is an angle small ([22]) if  $A = \bigcup_{n=1}^{\infty} A_n$ , where  $A_n$  are  $\alpha$ -angle porous.

Theorem 4 ([22]). Let X be a real Banach space such that  $X^{\pm}$  is separable,  $T: X \longrightarrow 2^{X^{\pm}}$  a monotone mapping with  $D(T) \subset X$ . Then there exists an angle small subset  $A \subset D(T)$  such that T is single-valued and upper semicontinuous ( with respect to the norm topologies of X and  $X^{\pm}$ ) on A.

Theorem 5 ([7]). Let X be a Banach space,  $f: X^{\mathbb{H}} \to \mathbb{R}$  a convex functional which is continuous with respect to the Mackey topology  $\mathcal{C}(X^{\mathbb{H}},X)$ . Then f is Fréchet differentiable on a norm-dense  $G_{\mathcal{F}}$  subset of  $X^{\mathbb{H}}$ .

According to  $\Gamma(18)$  a map  $f: X \to Y$  is said to be a Borel measurable function of the 1<sup>st</sup> Borel class if for each closed subset H of Y the set  $f^{-1}$  (H) is a  $G_f$  set in X.

Theorem 6 ([18]). Let X be a Banach space,  $T: X \longrightarrow 2^{X^{**}}$  a maximal monotone operator with int  $D(T) \ne \mathcal{O}$ . (i) If X admits an equivalent norm whose dual norm on  $X^{**}$  is (R), then T has a norm-to-weak Borel measurable selection  $T_0$  of the 1<sup>st</sup> Borel class on D(T). The set C of points of int D(T) where  $T_0$  is norm-to-weak  $T_0$ 

continuous coincides with the set of all points of int D(T) where  $T_o$  is point-valued. Further C contains a dense  $G_{\mathcal{F}}$  subset of int D(T). (ii) If  $X^{\mathbb{H}}$  has the Radon-Nikodým property, then T has a norm-to-norm measurable selection  $T_o$  of the 1<sup>St</sup> Borel class on D(T). The set U of all points of int D(T), at which  $T_o$  is norm-to-norm continuous, coincides with the set of all points of int D(T), at which T is point-valued and norm-to-norm upper semicontinuous. Furthermore U is dense  $G_{\mathcal{F}}$  subset of int D(T).

3. Accretive and maximal accretive multivalued mappings.

First of all we recall some basic and well-known notions concerning accretive operators. A multivalued mapping  $A: X \to 2^X$  is said to be: (i) accretive on D(A) if for each  $u, v \in D(A)$  and each  $x \in A(u)$ ,  $y \in A(v)$  there exists an element  $x^X \in J(u-v)$  such that  $\langle x-y, x^X \rangle \stackrel{\geq}{=} 0$ ; (ii) maximal accretive on D(A) if A is accretive on D(A) and if  $(u,x) \in X \times X$  is a given element such that for each  $v \in D(A)$  and  $y \in A(v)$  there exists  $x^X \in J(u-v)$  such that  $\langle x-y, x^X \rangle \stackrel{\geq}{=} 0$ , then  $u \in D(A)$  and  $x \in A(u)$ ;

(iii) hemicontinuous at  $u_0 \in \operatorname{int_aD}(A)$  (an algebraic interior of D(A)) if for each  $u \in X$ , every null sequence of positive numbers  $t_n$  and every  $v_n \in A(u_n)$ , where  $u_n = u_0 + t_n u$ ,  $(v_n)$  converges weakly in X to some point  $z_0 \in A(u_0)$ .

Theorem 7. Let X be a reflexive smooth and rotund Banach space, A:  $X^{\#} \rightarrow 2^{X^{\#}}$  an accretive mapping (with respect to the duality mapping J:  $X^{\#} \rightarrow X$ ) such that  $D(A) = X^{\#}$  and for each  $u^{\#} \in X^{\#}$   $A(u^{\#})$  is convex and closed in  $X^{\#}$ . If A is hemicontinuous on  $X^{\#}$ , then A is maximal accretive on  $X^{\#}$ .

Let us recall that Fabián [11] stated the following result: If X is a reflexive Banach space such that X,  $X^{X}$  are both (LUR) and A:  $X \rightarrow 2^{X}$  is maximal accretive such that int  $D(A) \neq \mathcal{X}$  and  $(A^{-1} + \lambda I)(X) = X$  for each  $\lambda > 0$ , then A is single-valued and upper semicontinuous ( with respect to the norm topology of X ) on a dense  $G_{P}$  subset of int D(A).

Theorem 8. Let X be a Banach space, A:  $X \rightarrow 2^X$  a maximal accretive mapping such that int D(A)  $\neq \mathcal{X}$ .

- (a) If X is reflexive and (F)-smooth, then there is a dense  $G_{\mathcal{F}}$  set  $D_{\mathcal{O}}$  int D(A) such that A|D is single-valued and continuous from the norm topology of X into the weak topology of X;
- (b) If X is (F)-smooth and the duality mapping  $J: X \rightarrow X^{M}$  is open, then A is single-valued and upper semicontinuous ( with respect to the norm topology of X ) on a dense  $G_{F}$  subset of int D(A).

Remark 1 . If X is reflexive smooth and (H)-Banach space, then J is open. In particular, if X is smooth and (LUR)-Banach space. then J is open. Note that if  $X^{H}$  is (LUR), then X is (F)-smooth and if X is reflexive and (LUR), then  $X^{X}$  is (F)-smooth. Since X and  $X^{X}$  are both (F)-smooth, J is a homeomorphism of X onto  $X^{*}$  ([16]).

Proposition 1 . If X is reflexive (F)-smooth Banach space, A:  $X \rightarrow 2^X$  is accretive on D(A) and lower semicontinuous at  $u_0 \in D(A)$ from the norm topology of X into the weak topology of X, then A(u,) is a singleton.

Theorem 8 and Proposition 1 show that the properties of maximal accretive multivalued mappings deeply rely on the structure of Banach spaces ( compare [19]) .

Theorem 9. Let X be a real normed linear space, f a convex continuous functional on X ,  $v_0$ ,  $w_0^{*}$  given points of X and  $X^{*}$ , respectively. Assume that there exists a closed linear subspace E of X such that  $\{u \in E : \mathcal{Y}_{v,w} * (u) \le c\}$  is non-empty and relatively weakly compact in E for some c > 0, where  $\mathcal{Y}$  is defined by  $\mathcal{Y}(u) = c$ = f (  $u + v_0$ ) -  $\langle w_0^{\Re}, u \rangle$  for each  $u \in E$ . Then :

(i) There exists a point 
$$u_0 \in E$$
 such that   
(\*) Of  $(u_0 + v_0) \cap (w_0^{\#} + E^{\perp}) \neq \theta$ .

(ii) If f is Gateaux differentiable at the point  $u_0 + v_0$ , then the intersection (\*) consists of exactly one point.

Corollary 1 . Let X be a real normed linear space, f a convex continuous functional on X . Assume that there exists a reflexive subspace E of X such that  $f(u) \cdot ||u||^{-1} \rightarrow +\infty$  as  $||u|| \rightarrow +\infty$ (i) If  $v_0, w_0^{\mathbb{H}}$  are arbitrary points of  $X, X^{\mathbb{H}}$  respectively, Then: then

$$\partial f'(u_0 + v_0) \wedge (w_0^* + E^{\perp}) \neq \emptyset$$
.

(ii) If f is Gateaux differentiable on X, then the above intersection consists of exactly one point.

Corollary 1 extends the results of Beurling and Livingston [3], Browder [5], Asplund [1]. Another generalization of the Beurling--Livingston theorem was given by Gobbo [17].

Further results concerning these topics will be published later.

#### REFERENCES

- [1] ASPLUND E. "Positivity of duality mappings", Bull. Amer. Math. Soc., 73 (1967), 200 203.
- 12] BARBU V. and PRECUPANU T. "Convexity and optimalization in Banach spaces", Ed. Academiei, Bucuresti, Sijthoff and Noordhoff, Int. Publ., 1978.
- [3] BEURLING A. and LIVINGSTON A.E. "A theorem on duality mapping in Banach spaces", Ark. Mat., 4 (1962), 405 411.
- E4] BROWDER F.E. "Nonlinear operators and nonlinear equations in Banach spaces", Proc. Sympos. Pure Math., 18, part 2, (1968), Amer. Math. Soc. Providence, 1976.
- [5] BROWDER F.E. "On a theorem of Beurling and Livingston", Canad. J. Math., 17 (1965), 367 372.
- CHRISTENSEN J.P.R. and KENDEROV P.S. "Dense strong continuity of mappings and the Radon-Nikodym property". Math. Scand., 54 (1984), 70 78.
- L71 CHRISTENSEN J.P.R. and KENDEROV P.S. "Dense Fréchet differentiability of Mackey continuous convex functions", Compt. Rend. Acad. Bulg. Sci., 36 (1983), 737 - 738.
- [8] DEIMLING KL. "Nonlinear functional analysis", Springer-Verlag, Berlin, 1985.
- [9] FABIÁN M. "On single-valuedness (and strong continuity) of maximal monotone mappings", Comment. Math. Univ. Carolinae, 18 (1977), 19 39.
- L10] FABIAN M. "Once more on continuity of maximal monotone mappings", Comment. Math. Univ. Carolinae, 18 (1977), 105 114.
- Lill FABIÁN M. "On strong continuity of monotone mappings in reflexive Banach spaces", Acta Pol., 6 (1977), 85 98.
- FITZPATRICK S.P. "Continuity of non-linear monotone operators", Proc. Amer. Math. Soc., 62 (1977), 111 116.
- Li31 FITZPATRICK S.P. "Monotone operators and dentability", Bul. Austr. Math. Soc. 18 (1978), 72 82.
- [14] FOWLER P.A. "Correction of a generalization of a theorem of Beurling and Livingston", Proc. Amer. Math. Soc., 74 (1979), 56 58.
- [15] GILES J.R. "Convex analysis with application in the differentiation of convex functions", Pitman, Melbourne, 1982.
- [16] GILES J.R. "A Banach space with support homeomorphism is reflexive", Bull. Austral. Math. Soc., 12 (1971), 393 396.

- [17] GOBBO L. "A generalization of a theorem of A. Beurling and A. E. Livingston", Boll. U. M. I. (6) 3 A (1984), 441 448.
- [18] JAYNE J.E. and ROGERS C.A. "Borel selectors for upper semicontinuous set-valued maps", Acta Math., 155 (1985), 41 - 79.
- (19] KENDEROV P.S. "Multivalued monotone mappings are almost everywhere single-valued", Studia Math., 56 (1976), 199 203.
- [20] KENDEROV P.S. "Monotone operators in Asplund spaces", Compt. rend. Acad. Sci. Bulgare, 30 (1977), 963 964.
- [21] PASCALI D. and SBURLAN "Nonlinear mappings of monotone type", Academiei Bucuresti, Sijthoff and Noordhoff, Int. Publ., 1978.
- PREISS D. and ZAJÍČEK L. "Stronger estimates of smallness of sets of Fréchet nondifferentiability of convex functions", Suppl. Rend. Circolo Mat. di Palermo, 1984, 219 223.
- [23] ZAJÍČEK L. "On the differentiation of convex functions in finite and infinite dimensional spaces", Czech. Math. J., 29 (1979), 340 363.
- [24] ZARANTONELLO E.H. "Dense single-valuedness of monotone operators", Israel J. of Math., 15 (1972), 158 166.

MATHEMATICAL INSTITUTE
OF CHARLES UNIVERSITY
SOKOLOVSKÁ 83
186 OO PRAHA 8 - KARLÍN