

Jiří Vanžura

The cohomology of $\tilde{G}_{m,2}$ with integer coefficients

In: Jan Slovák and Martin Čadek (eds.): Proceedings of the 18th Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1999. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 59. pp. 201–208.

Persistent URL: <http://dml.cz/dmlcz/702139>

Terms of use:

© Circolo Matematico di Palermo, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE COHOMOLOGY OF $\tilde{G}_{n,2}$ WITH INTEGER COEFFICIENTS

JIRÍ VANŽURA

ABSTRACT. This paper contains the description of the cohomology ring $H^*(\tilde{G}_{n,2}; \mathbb{Z})$ of the Grassmannian $\tilde{G}_{n,2}$ of oriented planes with integer coefficients. We describe the ring in terms of generators and relations.

1. INTRODUCTION AND PRELIMINARIES

The description of the cohomology ring $H^*(\tilde{G}_{n,2}; \mathbb{Z})$ seems not to be available in the literature. Its knowledge is necessary for example when studying the existence of 2-dimensional subbundles of a vector bundle. Another reason for writing this paper was the necessity to use some results from it in our forthcoming paper [CV] about the cohomology ring of the Grassmannian $G_{n,2}$ of nonoriented planes with integer and twisted integer coefficients.

We shall consider the vector space $\mathbb{R}^n$ with its canonical orientation. The symbol $\tilde{G}_{n,k}$, where $0 < k < n$, will denote the Grassmann manifold of oriented k -dimensional subspaces in $\mathbb{R}^n$, and $\tilde{\gamma}_k$ will be the canonical oriented k -dimensional vector bundle over $\tilde{G}_{n,k}$. The vector bundle $\tilde{\gamma}_k$ is obviously a subbundle of the trivial n -dimensional vector bundle ε^n with the fiber $\mathbb{R}^n$, and it has a riemannian metric induced from the canonical riemannian metric on ε^n . There is also the orthogonal complement $\tilde{\gamma}_k^\perp$, which is an $(n - k)$ -dimensional vector bundle. We orient it in such a way that the orientation of $\tilde{\gamma}_k \oplus \tilde{\gamma}_k^\perp$ coincides with the canonical orientation of ε^n .

From now on we shall assume that $n \geq 4$. We shall consider the unit sphere bundles

$$\pi_1 : S^1 \tilde{\gamma}_2 \longrightarrow \tilde{G}_{n,2}$$

and

$$\pi_2 : S^{n-2} \tilde{\gamma}_{n-1} \longrightarrow \tilde{G}_{n,n-1}.$$

1991 *Mathematics Subject Classification.* 55R40.

Key words and phrases. Grassmannian of oriented planes, Singular cohomology.

Research supported by the grant 201/96/0310 of the Grant Agency of the Czech Republic.

This paper is in final form and no version of it will be published elsewhere.

1. Lemma. *The total spaces $S^1\tilde{\gamma}_2$ and $S^{n-2}\tilde{\gamma}_{n-1}$ are homeomorphic.*

Proof. We can easily construct a mapping $\varphi : S^1\tilde{\gamma}_2 \rightarrow S^{n-2}\tilde{\gamma}_{n-1}$. An element from $S^1\tilde{\gamma}_2$ is a couple (α, v) , where $\alpha \subset \mathbb{R}^n$ is a 2-dimensional oriented subspace and $v \in \alpha$ is a unit vector. In α there is a unique unit vector v' orthogonal to v such that $\{v, v'\}$ is a positive basis in α . Further let $v^\perp$ denote the orthogonal complement to v endowed with such orientation that the natural orientation of $[v] \oplus v^\perp$ coincides with the orientation of $\mathbb{R}^n$. We can now define φ by the formula

$$\varphi(\alpha, v) = (v^\perp, v').$$

It is easy to verify that this mapping is a homeomorphism.

Our next aim is to compute the cohomology ring $H^*(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z})$. We shall use the Gysin sequence for the fibration

$$S^{n-2} \xrightarrow{i_3} S^{n-2}\tilde{\gamma}_{n-1} \xrightarrow{\pi_3} \tilde{G}_{n,n-1} \cong S^{n-1}.$$

We have to distinguish two cases.

2. n IS EVEN

The Euler class $e = e(\tilde{\gamma}_{n-1}) \in H^{n-1}(S^{n-1}; \mathbb{Z})$ has order two and consequently $e = 0$. Therefore we get the Gysin sequence in the form

$$\dots \xrightarrow{0} H^k(S^{n-1}) \xrightarrow{\pi_3^*} H^k(S^{n-2}\tilde{\gamma}_{n-1}) \rightarrow H^{k-n+2}(S^{n-1}) \xrightarrow{0} \dots$$

Thus we can immediately see that

$$H^k(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) = 0 \quad \text{for } k \neq 0, n-2, n-1, 2n-3.$$

Because $S^{n-2}\tilde{\gamma}_{n-1}$ is connected, we have

$$H^0(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) = \mathbb{Z}.$$

It is also easy to determine the remaining cohomology groups. For the later use we shall write here the relevant parts of the Gysin sequence. For $k = n-2$ we get

$$0 = H^{n-2}(S^{n-1}) \xrightarrow{\pi_3^*} H^{n-2}(S^{n-2}\tilde{\gamma}_{n-1}) \rightarrow H^0(S^{n-1}) = \mathbb{Z} \xrightarrow{0} \dots$$

which shows that

$$H^{n-2}(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) = \mathbb{Z}.$$

For $k = n-1$ we get

$$\dots \xrightarrow{0} \mathbb{Z} = H^{n-1}(S^{n-1}) \xrightarrow{\pi_3^*} H^{n-1}(S^{n-2}\tilde{\gamma}_{n-1}) \rightarrow H^1(S^{n-1}) = 0$$

which gives

$$H^{n-1}(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) = \mathbb{Z}.$$

Finally for $k = 2n - 3$ we have

$$0 = H^{2n-3}(S^{n-1}) \xrightarrow{\pi_2^*} H^{2n-3}(S^{n-2}\tilde{\gamma}_{n-1}) \rightarrow H^{n-1}(S^{n-1}) = \mathbb{Z} \xrightarrow{0} \dots$$

which gives

$$H^{2n-3}(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) = \mathbb{Z}.$$

Because $S^{n-2}\tilde{\gamma}_{n-1}$ is a manifold of dimension $2n - 3$ this determines the additive structure of the cohomology in question.

Now we shall describe generators in the above groups. Let us denote $\psi = \varphi^{-1} : S^{n-2}\tilde{\gamma}_{n-1} \rightarrow S^1\tilde{\gamma}_2$. It is easy to see that for an element $(\beta, w) \in S^{n-2}\tilde{\gamma}_{n-1}$, where $\beta \subset \mathbb{R}^n$ is an oriented $(n - 1)$ -dimensional subspace and $w \in \beta$ is a unit vector we get

$$\psi(\beta, w) = ([\beta^\perp, w], \beta^\perp),$$

where $\beta^\perp$ denotes the unique unit vector orthogonal to β and such that the natural orientation of $[\beta^\perp] \oplus \beta$ coincides with the orientation of $\mathbb{R}^n$. We orient the 2-dimensional subspace $[\beta^\perp, w]$ by taking $\{\beta^\perp, w\}$ as a positive basis. On $S^{n-2}\tilde{\gamma}_{n-1}$ we have an oriented $(n - 2)$ -dimensional vector bundle $\psi^*\pi_1^*\tilde{\gamma}_2^\perp$. Its fiber over the point (β, w) has the form

$$((\beta, w), w_\beta^\perp),$$

where $w_\beta^\perp$ denotes the $(n - 2)$ -dimensional subspace of β orthogonal to w and oriented in such a way that the natural orientation of $[w] \oplus w_\beta^\perp$ coincides with the orientation of β . If we consider again the fibration $S^{n-2} \xrightarrow{i_2} S^{n-2}\tilde{\gamma}_{n-1} \xrightarrow{\pi_2} S^{n-1}$ we can easily see that $i_2^*\psi^*\pi_1^*\tilde{\gamma}_2^\perp \cong TS^{n-2}$, where TS^{n-2} denotes the tangent bundle. Let us orient TS^{n-2} in such a way that this is an orientation preserving isomorphism. We denote $\omega \in H^{n-2}(S^{n-2}; \mathbb{Z})$ the generator uniquely determined by the orientation of TS^{n-2} . Obviously we have $e(i_2^*\psi^*\pi_1^*\tilde{\gamma}_2^\perp) = 2\omega$, where e denotes the Euler class. The Serre sequence for this fibration (with $\mathbb{Z}$ -coefficients) shows that i_2^* is an isomorphism. We denote by a the unique element $a \in H^{n-2}(S^{n-2}\tilde{\gamma}_{n-1})$ such that $i_2^*a = \omega$. (We shall also write a instead of φ^*a .) Obviously $a \in H^{n-2}(S^{n-2}\tilde{\gamma}_{n-1})$ is a generator. We have then

$$e(\psi^*\pi_1^*\tilde{\gamma}_2^\perp) = 2a.$$

The same Serre sequence shows also that $\pi_2^* : H^{n-1}(S^{n-1}) \rightarrow H^{n-1}(S^{n-2}\tilde{\gamma}_{n-1})$ is an isomorphism. Let $\theta \in H^{n-1}(S^{n-1})$ be a generator. We take a generator $b \in H^{n-1}(S^{n-2}\tilde{\gamma}_{n-1})$ such that $b = \pi_2^*\theta$.

We shall now prove that $c = ab$ is a generator of the group $H^{2n-3}(S^{n-2}\tilde{\gamma}_{n-1})$. For this purpose we shall use the oriented vector bundle $\tilde{\gamma}_{n-1}$ over S^{n-1} . Let

$$U \in H^{n-1}(\mathbb{B}^{n-1}\tilde{\gamma}_{n-1}, S^{n-2}\tilde{\gamma}_{n-1}).$$

(again with $\mathbb{Z}$ -coefficients) denote the Thom class. We then have

$$\delta a = \pm 1 \cup U.$$

Because $b = \pi_2^*\theta$ we have

$$\delta(ab) = \pm \theta \cup U,$$

which shows that ab is a generator. We have thus proved the following proposition.

2. Proposition. *The cohomology ring $H^*(S^{n-2}\tilde{\gamma}_{n-1}; \mathbf{Z})$ is isomorphic with the graded ring*

$$\mathbf{Z}[a, b]/(a^2, b^2),$$

where $\deg a = n - 2$, $\deg b = n - 1$.

We shall now use the above results for the computation of the cohomology ring $H^*(\tilde{G}_{n,2}; \mathbf{Z})$. (We recall that $n \geq 4$ is even.) For this purpose we shall use the Gysin sequence for the oriented vector bundle $\tilde{\gamma}_2$ over $\tilde{G}_{n,2}$. We shall denote $e = e(\tilde{\gamma}_2) \in H^2(\tilde{G}_{n,2}; \mathbf{Z})$ the Euler class of $\tilde{\gamma}_2$. First we shall prove the following proposition.

3. Proposition.

$$H^{2k}(\tilde{G}_{n,2}; \mathbf{Z}) \cong \mathbf{Z} \ni e^k \quad \text{for } 2k < n - 2$$

with e^k being a generator,

$$H^{2k+1}(\tilde{G}_{n,2}; \mathbf{Z}) = 0 \quad \text{for } 2k + 1 < n - 2.$$

Proof. Because $\tilde{G}_{n,2}$ is connected we have $H^0(\tilde{G}_{n,2}; \mathbf{Z}) \cong \mathbf{Z}$ with the generator $1 = e^0$. Because $\tilde{G}_{n,2}$ is simply connected we have $H^1(\tilde{G}_{n,2}; \mathbf{Z}) = 0$. Now it is sufficient to proceed by induction using the Gysin sequence for the vector bundle $\tilde{\gamma}_2$.

4. Proposition.

$$H^{2k}(\tilde{G}_{n,2}; \mathbf{Z}) \cong \mathbf{Z} \quad \text{for } 2n - 4 \geq 2k > n - 2,$$

$$H^{2k+1}(\tilde{G}_{n,2}; \mathbf{Z}) = 0 \quad \text{for } 2k + 1 > n - 2.$$

Proof. $\tilde{G}_{n,2}$ is an orientable compact manifold of dimension $2n - 4$. This implies that $H^{2n-4}(\tilde{G}_{n,2}; \mathbf{Z}) \cong \mathbf{Z}$. $\tilde{G}_{n,2}$ is simply connected, and consequently $H_1(\tilde{G}_{n,2}) = 0$. The Poincaré duality gives then $H^{2n-5}(\tilde{G}_{n,2}; \mathbf{Z}) = 0$. Now, it is again sufficient to proceed by induction (going down) using the same Gysin sequence as above.

It remains to compute the group $H^{n-2}(\tilde{G}_{n,2}; \mathbf{Z})$. The relevant part of the Gysin sequence has the form

$$0 = H^{n-3}(S^1\tilde{\gamma}_2) \rightarrow H^{n-4}(\tilde{G}_{n,2}) = \mathbf{Z} \xrightarrow{\cup e} H^{n-2}(\tilde{G}_{n,2}) \xrightarrow{\pi_1^*}$$

$$\xrightarrow{\pi_1^*} H^{n-2}(S^1\tilde{\gamma}_2) = \mathbf{Z} \rightarrow H^{n-3}(\tilde{G}_{n,2}) = 0,$$

which gives

$$H^{n-2}(\tilde{G}_{n,2}; \mathbf{Z}) \cong \mathbf{Z} \oplus \mathbf{Z}.$$

It is easy to see that in this group we can choose the generators $e^{(n-2)/2}$ and f' , where f' satisfies $\pi_1^* f' = a$. Obviously the Euler class $e(\tilde{\gamma}_2^\perp)$ can be uniquely expressed in the form

$$e(\tilde{\gamma}_2^\perp) = ue^{(n-2)/2} + 2f'.$$

We shall use the standard notation $w_i = w_i(\tilde{\gamma}_2)$ for the Stiefel-Whitney classes. From the relation $\tilde{\gamma}_2 \oplus \tilde{\gamma}_2^\perp = \epsilon^n$, where ϵ^n denotes the trivial n -dimensional vector bundle, we can easily find that $w_2(\tilde{\gamma}_2^\perp) = w_2$. This shows that the integer u is odd, i. e. $u = 2v + 1$. Taking $f = ve^{(n-2)/2} + f'$ we obtain for $H^{n-2}(\tilde{G}_{n,2}; \mathbb{Z})$ the generators $e^{(n-2)/2}$ and f , and for the Euler class $e(\tilde{\gamma}_2^\perp)$ we have the formula

$$e(\tilde{\gamma}_2^\perp) = e^{(n-2)/2} + 2f.$$

Moreover the same relation $\tilde{\gamma}_2 \oplus \tilde{\gamma}_2^\perp = \epsilon^n$ shows that

$$e^{n/2} = -2ef.$$

Let us consider now the following part of our Gysin sequence.

$$\mathbb{Z} \oplus \mathbb{Z} = H^{n-2}(\tilde{G}_{n,2}) \xrightarrow{\cup e} H^n(\tilde{G}_{n,2}) = \mathbb{Z} \rightarrow H^n(S^1\tilde{\gamma}_2) = 0$$

We can see that $e^{(n-2)/2} + 2f$ and f is a base of $H^{n-2}(\tilde{G}_{n,2})$ considered as a free module over $\mathbb{Z}$. Because $(e^{(n-2)/2} + 2f) \cup e = 0$ it is obvious that ef is a generator. We have

$$H^{n-2}(\tilde{G}_{n,2}; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z} \text{ with generators } e^{(n-2)/2} \text{ and } f.$$

Using this result and the same Gysin sequence as above we obtain by induction

$$H^{2k}(\tilde{G}_{n,2}; \mathbb{Z}) \cong \mathbb{Z} \text{ with generator } e^{(2k-n+2)/2}f \text{ for } 2n-4 \geq 2k > n-2.$$

It remains to determine f^2 . Obviously $f^2 = te^{(n-2)/2}f$, where t is an integer. Our next aim is to determine this integer. We shall apply the Gysin sequence of the vector bundle

$$\pi^\perp : \tilde{\gamma}_2^\perp \rightarrow \tilde{G}_{n,2}$$

with $\mathbb{Z}$ -coefficients. More precisely, we shall need only the following piece of the Gysin sequence.

$$H^{n-2}(\tilde{G}_{n,2}) \xrightarrow{\cup e^\perp} H^{2n-4}(\tilde{G}_{n,2}) \rightarrow H^{2n-4}(S^{n-3}\tilde{\gamma}_2^\perp),$$

where $S^{n-3}\tilde{\gamma}_2^\perp$ is the sphere bundle of $\tilde{\gamma}_2^\perp$ and $e^\perp = e(\tilde{\gamma}_2^\perp)$. We know already that $e^\perp = e^{(n-2)/2} + 2f$.

Obviously, we must first calculate $H^{2n-4}(S^{n-3}\tilde{\gamma}_2^\perp)$. Let us notice first that $S^{n-3}\tilde{\gamma}_2^\perp$ is a compact connected orientable manifold and $\dim S^{n-3}\tilde{\gamma}_2^\perp = 3n-7$. For this purpose we shall consider the following part of the Gysin sequence for the vector bundle $\tilde{\gamma}_2^\perp$.

$$\begin{aligned} 0 &= H^{n-3}(\tilde{G}_{n,2}) \rightarrow H^{n-3}(S^{n-3}\tilde{\gamma}_2^\perp) \rightarrow H^0(\tilde{G}_{n,2}) = \mathbb{Z} \xrightarrow{\cup e^\perp} \\ &\xrightarrow{\cup e^\perp} H^{n-2}(\tilde{G}_{n,2}) = \mathbb{Z} \oplus \mathbb{Z} \rightarrow H^{n-2}(S^{n-3}\tilde{\gamma}_2^\perp) \rightarrow H^1(\tilde{G}_{n,2}) = 0 \end{aligned}$$

From this sequence we get immediately

$$H^{n-3}(S^{n-3}\tilde{\gamma}_2^\perp) = 0$$

$$H^{n-2}(S^{n-3}\tilde{\gamma}_2^\perp) = \mathbf{Z} \quad \text{with the generator } (\pi^\perp)^* f.$$

Using the Poincare duality on $S^{n-3}\tilde{\gamma}_2^\perp$ we get

$$H_{2n-5}(S^{n-3}\tilde{\gamma}_2^\perp) = \mathbf{Z}, \quad H_{2n-4}(S^{n-3}\tilde{\gamma}_2^\perp) = 0.$$

Now it is sufficient to apply the universal coefficients theorem. We can write the exact sequence

$$\begin{aligned} 0 \rightarrow \text{Ext}(H_{2n-5}(S^{n-3}\tilde{\gamma}_2^\perp), \mathbf{Z}) \rightarrow H^{2n-4}(S^{n-3}\tilde{\gamma}_2^\perp; \mathbf{Z}) \rightarrow \\ \rightarrow \text{Hom}(H_{2n-4}(S^{n-3}\tilde{\gamma}_2^\perp), \mathbf{Z}) \rightarrow 0, \end{aligned}$$

which shows that

$$H^{2n-4}(S^{n-3}\tilde{\gamma}_2^\perp; \mathbf{Z}) = 0.$$

Now we can see that the mapping

$$\cup e^\perp : H^{n-2}(\tilde{G}_{n,2}) \rightarrow H^{2n-4}(\tilde{G}_{n,2})$$

is surjective. We have

$$\begin{aligned} (e^{(n-2)/2} + 2f)(e^{(n-2)/2} + 2f) &= 2e^{(n-2)/2}f + 4f^2 = (2 + 4t)e^{(n-2)/2}f \\ f(e^{(n-2)/2} + 2f) &= e^{(n-2)/2}f + 2f^2 = (1 + 2t)e^{(n-2)/2}f \end{aligned}$$

Consequently, there exists integers r and s such that

$$r(2 + 4t) + s(1 + 2t) = 1.$$

This equation can be written in the form

$$(1 + 2t)(2r + s) = 1,$$

which shows that $t = 0$. We have thus proved that

$$f^2 = 0.$$

We have obtained in this way a description of the integral cohomology ring of $\tilde{G}_{n,2}$.

5. Theorem. *The cohomology ring $H^*(\tilde{G}_{n,2}; \mathbf{Z})$ with n even, $n \geq 4$ is isomorphic with the graded ring*

$$\mathbf{Z}[e, f]/(e^{n/2} + 2ef, f^2),$$

where $\deg e = 2$, $\deg f = n - 2$. Moreover, under this isomorphism, $e(\tilde{\gamma}_2^\perp)$ correspond to the class determined by the element $e^{(n-2)/2} + 2f$.

3. n IS ODD

Let us denote by $\omega \in H^{n-1}(S^{n-1}; \mathbb{Z})$ a generator of S^{n-1} . Because $\tilde{\gamma}_{n-1}$ is isomorphic with TS^{n-1} we have $e = e(\tilde{\gamma}_{n-1}) = \pm 2\omega$. Obviously, we can choose ω in such a way that $e = e(\tilde{\gamma}_{n-1}) = 2\omega$. The same Gysin sequence as in the even case gives us first

$$H^k(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) = 0 \quad \text{for } k \neq 0, n-2, n-1, 2n-3.$$

We have obviously

$$H^0(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) \cong \mathbb{Z}.$$

For $k = n-2$ we get

$$0 = H^{n-2}(S^{n-1}) \rightarrow H^{n-2}(S^{n-2}\tilde{\gamma}_{n-1}) \rightarrow H^0(S^{n-1}) = \mathbb{Z} \xrightarrow{\cup 2\omega} H^{n-1}(S^{n-1}) = \mathbb{Z},$$

which shows that

$$H^{n-2}(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) = 0.$$

Next for $k = n-1$ we have

$$\mathbb{Z} = H^0(S^{n-1}) \xrightarrow{\cup 2\omega} H^{n-1}(S^{n-1}) = \mathbb{Z} \rightarrow H^{n-1}(S^{n-2}\tilde{\gamma}_{n-1}) \rightarrow H^1(S^{n-1}) = 0,$$

which gives

$$H^{n-1}(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) \cong \mathbb{Z}_2.$$

Finally for $k = 2n-3$ we get

$$0 = H^{2n-3}(S^{n-1}) \rightarrow H^{2n-3}(S^{n-2}\tilde{\gamma}_{n-1}) \rightarrow H^{n-1}(S^{n-1}) = \mathbb{Z} \rightarrow H^{2n-2}(S^{n-1}) = 0,$$

which gives

$$H^{2n-3}(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z}) \cong \mathbb{Z}.$$

The following proposition is obvious.

6. Proposition. *The cohomology ring $H^*(S^{n-2}\tilde{\gamma}_{n-1}; \mathbb{Z})$ is isomorphic with the graded ring*

$$\mathbb{Z}[x, y]/(2x, x^2, xy, y^2),$$

where $\deg x = n-1$ and $\deg y = 2n-3$.

We shall now again compute the integral cohomology ring of $\tilde{G}_{n,2}$. We start with the following proposition.

7. Proposition.

$$H^i(\tilde{G}_{n,2}; \mathbb{Z}) = 0 \quad \text{for } i \text{ odd.}$$

Proof. $\tilde{G}_{n,2}$ is simply connected. Therefore $H_1(\tilde{G}_{n,2}; \mathbb{Z}) = 0$ and $H^1(\tilde{G}_{n,2}; \mathbb{Z}) = 0$. The Poincaré duality gives $H^{2n-5}(\tilde{G}_{n,2}; \mathbb{Z}) = 0$. Proceeding now by induction (first going up, then going down), and using the Gysin sequence for the vector bundle $\tilde{\gamma}_2$, we get easily the assertion.

8. Proposition.

$$H^{2k}(\tilde{G}_{n,2}; \mathbb{Z}) \cong \mathbb{Z} \quad \text{for } 2k < n - 1.$$

with the generator being e^k .

Proof. Obviously $H^0(\tilde{G}_{n,2}; \mathbb{Z}) \cong \mathbb{Z}$ with the generator $1 = e^0$. Now, it suffices to proceed by induction (going up) and use the same Gysin sequence as above.

9. Proposition.

$$H^{2k}(\tilde{G}_{n,2}; \mathbb{Z}) \cong \mathbb{Z} \quad \text{for } n - 1 < 2k \leq 2n - 4.$$

Proof. We proceed in the same way as above with the induction going down.

It remains to determine the group $H^{n-1}(\tilde{G}_{n,2}; \mathbb{Z})$. The Gysin sequence gives here

$$\begin{aligned} 0 = H^{n-2}(S^1\tilde{\gamma}_2) \rightarrow H^{n-3}(\tilde{G}_{n,2}) = \mathbb{Z} \xrightarrow{\cup e} H^{n-1}(\tilde{G}_{n,2}) \rightarrow \\ \rightarrow H^{n-1}(S^1\tilde{\gamma}_2) = \mathbb{Z}_2 \rightarrow H^{n-2}(\tilde{G}_{n,2}) = 0. \end{aligned}$$

The last group vanishes because $n - 2$ is odd. From this exact sequence we can see that $H^{n-1}(\tilde{G}_{n,2}) \cong \mathbb{Z}$ or $\mathbb{Z} \oplus \mathbb{Z}_2$. Associated with the exact coefficient sequence $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}_2 \rightarrow 0$ we have the exact sequence

$$0 = H^{n-2}(\tilde{G}_{n,2}; \mathbb{Z}_2) \xrightarrow{\beta} H^{n-1}(\tilde{G}_{n,2}; \mathbb{Z}) \xrightarrow{2\chi} H^{n-1}(\tilde{G}_{n,2}; \mathbb{Z}),$$

where β is the Bockstein homomorphism. We can see that the homomorphism 2χ is injective and consequently $H^{n-1}(\tilde{G}_{n,2}; \mathbb{Z}) \cong \mathbb{Z}$. We have the following exact sequence.

$$\begin{aligned} 0 = H^{n-2}(S^1\tilde{\gamma}_2) \rightarrow H^{n-3}(\tilde{G}_{n,2}) = \mathbb{Z} \xrightarrow{\cup e} H^{n-1}(\tilde{G}_{n,2}) = \mathbb{Z} \rightarrow \\ \rightarrow H^{n-1}(S^1\tilde{\gamma}_2) = \mathbb{Z}_2 \rightarrow H^{n-2}(\tilde{G}_{n,2}) = 0 \end{aligned}$$

Now it is obvious that we can choose a generator $f \in H^{n-1}(\tilde{G}_{n,2})$ in such a way that there is

$$e^{(n-1)/2} = 2f.$$

10. Proposition.

$$H^{2k}(\tilde{G}_{n,2}; \mathbb{Z}) \cong \mathbb{Z} \ni e^{(2k-n+1)/2} f \quad \text{for } n - 1 \leq 2k \leq 2n - 4.$$

with $e^{(2k-n+1)/2} f$ being a generator.

Proof. It is of the same type as before.

Now we get easily

11. Theorem. *The cohomology ring $H^*(\tilde{G}_{n,2}; \mathbb{Z})$ with n odd, $n \geq 5$ is isomorphic with the graded ring*

$$\mathbb{Z}[e, f] / (e^{(n-1)/2} - 2f, f^2),$$

where $\deg e = 2$, $\deg f = n - 1$. Moreover, there is $e(\tilde{\gamma}_2^\perp) = 0$.

REFERENCES

[CV] M. Čadek, J. Vanžura, *The cohomology of $G_{m,2}$ with twisted integer coefficients*, to appear.

INSTITUTE OF MATHEMATICS, ACADEMY OF SCIENCES, ŽITKOVA 22, 616 69 BRNO, CZECH REPUBLIC

E-mail address: vanzura@ipm.cz